

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Remodulin safely and effectively. See full prescribing information for Remodulin.

**REMODULIN (treprostinil) Injection**  
Initial U.S. Approval: May 2002

### RECENT MAJOR CHANGES

Dosage and Administration (2.1) 01/2010  
Warnings and Precautions (5.1) 01/2010

### INDICATIONS AND USAGE

Remodulin is a prostacyclin vasodilator indicated for:

- Treatment of pulmonary arterial hypertension (PAH) (WHO Group 1) to diminish symptoms associated with exercise. Studies establishing effectiveness included patients with NYHA Functional Class II-IV symptoms and etiologies of idiopathic or heritable PAH (58%), PAH associated with congenital systemic-to-pulmonary shunts (23%), or PAH associated with connective tissue diseases (19%) (1.1)
- Patients who require transition from Flolan, to reduce the rate of clinical deterioration. The risks and benefits of each drug should be carefully considered prior to transition. (1.2)

### DOSAGE AND ADMINISTRATION

PAH in patients with NYHA Class II-IV symptoms:

- Initial dose for patients new to prostacyclin infusion therapy: 1.25 ng/kg/min (or 0.625 ng/kg/min if not tolerated); dose increase based on clinical response (increments of 1.25 ng/kg/min per week for the first 4 weeks of treatment, later 2.5 ng/kg/min per week). Limited experience with doses >40 ng/kg/min. Abrupt cessation of infusion should be avoided. (2.2, 2.3)
- Mild to moderate hepatic insufficiency: Initial dose should be decreased to 0.625 ng/kg/min ideal body weight; cautious dosage increase. Severe hepatic insufficiency: No studies performed. (2.4)

### Transition from Flolan:

Increase the Remodulin dose gradually as the Flolan dose is decreased, based on constant observation of response. (2.7)

### Administration:

Continuous subcutaneous infusion (undiluted) is the preferred mode. Use intravenous infusion (dilution required) if subcutaneous infusion is not tolerated. (2.1, 2.6)  
See Full Prescribing Information.

### DOSAGE FORMS AND STRENGTHS

- Remodulin is supplied in 20 mL vials containing 20, 50, 100, or 200 mg of treprostinil (1 mg/mL, 2.5 mg/mL, 5 mg/mL or 10 mg/mL). (3)

### CONTRAINDICATIONS

None

### WARNINGS AND PRECAUTIONS

- Chronic intravenous infusions of Remodulin are delivered using an indwelling central venous catheter. This route is associated with the risk of blood stream infections (BSIs) and sepsis, which may be fatal. (5.1)
- Remodulin should be used only by clinicians experienced in the diagnosis and treatment of PAH. (5.2)
- Adjust dosage based on clinical response, including infusion site symptoms. (5.3)
- Do not abruptly lower the dose or withdraw dosing. (5.4)

### ADVERSE REACTIONS

Most common adverse reactions (incidence >3%) reported in clinical studies with Remodulin: subcutaneous infusion site pain and reaction, headache, diarrhea, nausea, jaw pain, vasodilatation, dizziness, edema, pruritus and hypotension. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact United Therapeutics Corp. at 1-866-458-6479 or via e-mail at [drugsafety@unither.com](mailto:drugsafety@unither.com), or contact FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

### DRUG INTERACTIONS

- Blood pressure lowering drugs (e.g., diuretics, antihypertensive agents, or vasodilators): Risk of increased reduction in blood pressure (7.1)
- Remodulin inhibits platelet aggregation. Potential for increased risk of bleeding, particularly among patients on anticoagulants. (7.2)
- Remodulin dosage adjustment may be necessary if inhibitors or inducers of CYP2C8 are added or withdrawn. (7.6)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 2/2011

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### 1. INDICATIONS AND USAGE

- [1.1 Pulmonary Arterial Hypertension](#)
- [1.2 Pulmonary Arterial Hypertension in Patients Requiring Transition from Flolan®](#)

### 2. DOSAGE AND ADMINISTRATION

- [2.1 General](#)
- [2.2 Initial Dose for Patients New to Prostacyclin Infusion Therapy](#)
- [2.3 Dosage Adjustments](#)
- [2.4 Patients with Hepatic Insufficiency](#)
- [2.5 Patients with Renal Insufficiency](#)
- [2.6 Administration](#)
- [2.7 Patients Requiring Transition from Flolan](#)

### 3. DOSAGE FORMS AND STRENGTHS

### 4. CONTRAINDICATIONS

### 5. WARNINGS AND PRECAUTIONS

- [5.1 Risks Attributable to the Drug Delivery System](#)
- [5.2 General Conditions of Use](#)
- [5.3 Dose Modification](#)
- [5.4 Abrupt Withdrawal or Sudden Large Dose Reduction](#)
- [5.5 Patients with Hepatic or Renal Insufficiency](#)
- [5.6 Effect of Other Drugs on Treprostinil](#)

### 6. ADVERSE REACTIONS

- [6.1 Clinical Trials Experience](#)
- [6.2 Post-Marketing Experience](#)

### 7. DRUG INTERACTIONS

- [7.1 Antihypertensive Agents or Other Vasodilators](#)

### [7.4 Sildenafil](#)

### [7.5 Effect of Treprostinil on Cytochrome P450 Enzymes](#)

### [7.6 Effect of Cytochrome P450 Inhibitors and Inducers on Treprostinil](#)

### [7.7 Effect of Other Drugs on Treprostinil](#)

### 8. USE IN SPECIFIC POPULATIONS

- [8.1 Pregnancy](#)
- [8.2 Labor and Delivery](#)
- [8.3 Nursing Mothers](#)
- [8.4 Pediatric Use](#)
- [8.5 Geriatric Use](#)
- [8.6 Patients with Hepatic Insufficiency](#)
- [8.7 Patients with Renal Insufficiency](#)

### 10. OVERDOSAGE

### 11. DESCRIPTION

### 12. CLINICAL PHARMACOLOGY

- [12.1 Mechanism of Action](#)
- [12.2 Pharmacodynamics](#)
- [12.3 Pharmacokinetics](#)

### 13. NONCLINICAL TOXICOLOGY

- [13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility](#)

### 14. CLINICAL STUDIES

- [14.1 Clinical Trials in Pulmonary Arterial Hypertension \(PAH\)](#)
- [14.2 Flolan-To-Remodulin Transition Study](#)

### 16. HOW SUPPLIED / STORAGE AND HANDLING

### 17. PATIENT COUNSELING INFORMATION

\*Sections or subsections omitted from the full prescribing information are not listed.

## FULL PRESCRIBING INFORMATION

### REMODULIN® (treprostinil) Injection

#### 1. INDICATIONS AND USAGE

##### 1.1 Pulmonary Arterial Hypertension

Remodulin is indicated for the treatment of pulmonary arterial hypertension (PAH) (WHO Group 1) to diminish symptoms associated with exercise. Studies establishing effectiveness included patients with NYHA Functional Class II-IV symptoms and etiologies of idiopathic or heritable PAH (58%), PAH associated with congenital systemic-to-pulmonary shunts (23%), or PAH associated with connective tissue diseases (19%) [see *Clinical Studies (14.1)*].

It may be administered as a continuous subcutaneous infusion or continuous intravenous infusion; however, because of the risks associated with chronic indwelling central venous catheters, including serious blood stream infections, continuous intravenous infusion should be reserved for patients who are intolerant of the subcutaneous route, or in whom these risks are considered warranted.

##### 1.2 Pulmonary Arterial Hypertension in Patients Requiring Transition from Flolan®

In patients with pulmonary arterial hypertension requiring transition from Flolan (epoprostenol sodium), Remodulin is indicated to diminish the rate of clinical deterioration. The risks and benefits of each drug should be carefully considered prior to transition.

#### 2 DOSAGE AND ADMINISTRATION

##### 2.1 General

Remodulin is supplied in 20 mL vials containing 20, 50, 100, or 200 mg of treprostinil (1 mg/mL, 2.5 mg/mL, 5 mg/mL or 10 mg/mL). Remodulin can be administered as supplied or diluted for intravenous infusion with Sterile Water for Injection, 0.9% Sodium Chloride Injection, or Flolan® Sterile Diluent for Injection prior to administration.

##### 2.2 Initial Dose for Patients New to Prostacyclin Infusion Therapy

Remodulin is indicated for subcutaneous (SC) or intravenous (IV) use only as a continuous infusion. Remodulin is preferably infused subcutaneously, but can be administered by a central intravenous line if the subcutaneous route is not tolerated, because of severe site pain or reaction. The infusion rate is initiated at 1.25 ng/kg/min. If this initial dose cannot be tolerated because of systemic effects, the infusion rate should be reduced to 0.625 ng/kg/min.

##### 2.3 Dosage Adjustments

The goal of chronic dosage adjustments is to establish a dose at which PAH symptoms are improved, while minimizing excessive pharmacologic effects of Remodulin (headache, nausea, emesis, restlessness, anxiety and infusion site pain or reaction).

The infusion rate should be increased in increments of 1.25 ng/kg/min per week for the first four weeks of treatment and then 2.5 ng/kg/min per week for the remaining duration of infusion, depending on clinical response. Dosage adjustments may be undertaken more often if tolerated. There is little experience with doses >40 ng/kg/min. Abrupt cessation of infusion should be

avoided [see Warnings and Precautions (5.4)]. Restarting a Remodulin infusion within a few hours after an interruption can be done using the same dose rate. Interruptions for longer periods may require the dose of Remodulin to be re-titrated.

## 2.4 Patients with Hepatic Insufficiency

In patients with mild or moderate hepatic insufficiency, the initial dose of Remodulin should be decreased to 0.625 ng/kg/min ideal body weight and should be increased cautiously. Remodulin has not been studied in patients with severe hepatic insufficiency [see Warnings and Precautions (5.5), Use In Specific Populations (8.6) and Clinical Pharmacology (12.3)].

## 2.5 Patients with Renal Insufficiency

No studies have been performed in patients with renal insufficiency. No specific advice about dosing in patients with renal impairment can be given. [see Clinical Pharmacology (12.3)].

## 2.6 Administration

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit. If either particulate matter or discoloration is noted, Remodulin should not be administered.

### Subcutaneous Infusion

Remodulin is administered subcutaneously by continuous infusion, via a self-inserted subcutaneous catheter, using an infusion pump designed for subcutaneous drug delivery. To avoid potential interruptions in drug delivery, the patient must have immediate access to a backup infusion pump and subcutaneous infusion sets. The ambulatory infusion pump used to administer Remodulin should: (1) be small and lightweight, (2) be adjustable to approximately 0.002 mL/hr, (3) have occlusion/no delivery, low battery, programming error and motor malfunction alarms, (4) have delivery accuracy of  $\pm 6\%$  or better and (5) be positive pressure driven. The reservoir should be made of polyvinyl chloride, polypropylene or glass.

For subcutaneous infusion, Remodulin is **delivered without further dilution** at a calculated Subcutaneous Infusion Rate (mL/hr) based on a patients Dose (ng/kg/min), Weight (kg), and the Vial Strength (mg/mL) of Remodulin being used. During use, a single reservoir (syringe) of undiluted Remodulin can be administered up to 72 hours at 37°C. The Subcutaneous Infusion rate is calculated using the following formula:

$$\text{Subcutaneous Infusion Rate (mL/hr)} = \frac{\text{Dose (ng/kg/min)} \times \text{Weight (kg)} \times 0.00006^*}{\text{Remodulin Vial Strength (mg/mL)}}$$

\*Conversion factor of 0.00006 = 60 min/hour x 0.000001 mg/ng

Example calculations for **Subcutaneous Infusion** are as follows:

**Example 1:**

For a 60 kg person at the recommended initial dose of 1.25 ng/kg/min using the 1 mg/mL Remodulin Vial Strength, the infusion rate would be calculated as follows:

$$\text{Subcutaneous Infusion Rate (mL/hr)} = \frac{1.25 \text{ ng/kg/min} \times 60 \text{ kg} \times 0.00006}{1 \text{ mg/mL}} = 0.005 \text{ mL/hr}$$

**Example 2:**

For a 65 kg person at a dose of 40 ng/kg/min using the 5 mg/mL Remodulin Vial Strength, the infusion rate would be calculated as follows:

$$\text{Subcutaneous Infusion Rate (mL/hr)} = \frac{40 \text{ ng/kg/min} \times 65 \text{ kg} \times 0.00006}{5 \text{ mg/mL}} = 0.031 \text{ mL/hr}$$

**Intravenous Infusion**

**Remodulin must be diluted with either Sterile Water for Injection, 0.9% Sodium Chloride Injection, or Flolan® Sterile Diluent for Injection** and is administered intravenously by continuous infusion, via a surgically placed indwelling central venous catheter, using an infusion pump designed for intravenous drug delivery. If clinically necessary, a temporary peripheral intravenous cannula, preferably placed in a large vein, may be used for short term administration of Remodulin. Use of a peripheral intravenous infusion for more than a few hours may be associated with an increased risk of thrombophlebitis. To avoid potential interruptions in drug delivery, the patient must have immediate access to a backup infusion pump and infusion sets. The ambulatory infusion pump used to administer Remodulin should: (1) be small and lightweight, (2) have occlusion/no delivery, low battery, programming error and motor malfunction alarms, (3) have delivery accuracy of ±6% or better of the hourly dose, and (4) be positive pressure driven. The reservoir should be made of polyvinyl chloride, polypropylene or glass.

Diluted Remodulin has been shown to be stable at ambient temperature for up to 48 hours at concentrations as low as 0.004 mg/mL (4,000 ng/mL).

When using an appropriate infusion pump and reservoir, a predetermined intravenous infusion rate should first be selected to allow for a desired infusion period length of up to 48 hours between system changeovers. Typical intravenous infusion system reservoirs have volumes of 50 or 100 mL. With this selected Intravenous Infusion Rate (mL/hr) and the patient's Dose (ng/kg/min) and Weight (kg), the Diluted Intravenous Remodulin Concentration (mg/mL) can be calculated using the following formula:

**Step 1**

$$\text{Diluted Intravenous Remodulin Concentration (mg/mL)} = \frac{\text{Dose (ng/kg/min)} \times \text{Weight (kg)} \times 0.00006}{\text{Intravenous Infusion Rate (mL/hr)}}$$

The Amount of Remodulin Injection needed to make the required Diluted Intravenous Remodulin Concentration for the given reservoir size can then be calculated using the following formula:

**Step 2**

$$\text{Amount of Remodulin Injection (mL)} = \frac{\text{Diluted Intravenous Remodulin Concentration (mg/mL)}}{\text{Remodulin Vial Strength (mg/mL)}} \times \text{Total Volume of Diluted Remodulin Solution in Reservoir (mL)}$$

The calculated amount of Remodulin Injection is then added to the reservoir along with the sufficient volume of diluent (Sterile Water for Injection, 0.9% Sodium Chloride Injection, or Flolan® Sterile Diluent for Injection) to achieve the desired total volume in the reservoir.

Example calculations for ***Intravenous Infusion*** are as follows:

**Example 3:**

For a 60 kg person at a dose of 5 ng/kg/min, with a predetermined intravenous infusion rate of 1 mL/hr and a reservoir of 50 mL, the Diluted Intravenous Remodulin Concentration would be calculated as follows:

**Step 1**

$$\text{Diluted Intravenous Remodulin Concentration (mg/mL)} = \frac{5 \text{ ng/kg/min} \times 60 \text{ kg} \times 0.00006}{1 \text{ mL/hr}} = 0.018 \text{ mg/mL (18,000 ng/mL)}$$

The Amount of Remodulin Injection (using 1 mg/mL Vial Strength) needed for a total Diluted Remodulin Concentration of 0.018 mg/mL and a total volume of 50 mL would be calculated as follows:

**Step 2**

$$\text{Amount of Remodulin Injection (mL)} = \frac{0.018 \text{ mg/mL}}{1 \text{ mg/mL}} \times 50 \text{ mL} = 0.9 \text{ mL}$$

The Diluted Intravenous Remodulin Concentration for the person in Example 3 would thus be prepared by adding 0.9 mL of 1 mg/mL Remodulin Injection to a suitable reservoir along with a sufficient volume of diluent to achieve a total volume of 50 mL in the reservoir. The pump flow rate for this example would be set at 1 mL/hr.

**Example 4:**

For a 75 kg person at a dose of 30 ng/kg/min, with a predetermined intravenous infusion rate of 2 mL/hr, and a reservoir of 100 mL, the Diluted Intravenous Remodulin Concentration would be calculated as follows:

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.