

**Cited by (CrossRef):**

4 articles

[Check for updates](#)[Citation tools](#)

S. Kuramitsu, Department of Biology, Graduate School of Science, Osaka University, 1-1 Machikaneyama-cho, Toyonaka, Osaka  
+ 81 6 850 5433. Fax: + 81 6 850 5442.

E-mail: [kuramitu@bio.sci.osaka-u.ac.jp](mailto:kuramitu@bio.sci.osaka-u.ac.jp)

## Abstract

Thermostable RecA protein (ttRecA) from *Thermus thermophilus* HB8 showed strand exchange activity at 65 °C but not at 37 °C. A nucleoprotein complex was observed at both temperatures. ttRecA showed single-stranded DNA (ssDNA)-dependent ATPase activity. ATPase activity was maximal at 65 °C. The kinetic parameters,  $K_m$  and  $k_{cat}$ , for adenosine triphosphate (ATP) hydrolysis with poly(dA) were 0.60 s<sup>-1</sup> at 65 °C, and 0.34 mM and 0.28 s<sup>-1</sup> at 37 °C, respectively. Substrate cooperativity was observed at both temperatures. The Hill coefficient was about 2. At 65 °C, all tested ssDNAs were able to stimulate the ATPase activity. The order of ATPase stimulation was poly(dC) > poly(dT) > M13 ssDNA > poly(dA). Double-stranded DNAs (dsDNA), poly(dT)-poly(dA) and M13 dsDNA, were not stimulated by the enzyme at 65 °C. At 37 °C, however, not only dsDNAs but also poly(dA) and M13 ssDNA showed poor stimulating ability. M13 ssDNA gave circular dichroism (CD) peaks at around 192 nm, which reflect a particular structure of DNA. The conformation of ssDNA was altered by an upshift of temperature or binding to *Escherichia coli* RecA protein (ecRecA), but not to ttRecA. The dissociation constant of ecRecA and poly(dA) was estimated to be 44 μM at 25 °C by the change in the CD. These observations suggest that the capability of ssDNA to induce the conformation of ssDNA may be different between ttRecA and ecRecA. The specific structure of ssDNA was altered by heat treatment. After this alteration, ttRecA and ecRecA can express their activities at each physiological temperature.

## Abbreviations

**ATP-γ-S**

adenosine 5'-O-(3-thiotriphosphate)

**CD**

circular dichroism

**ecRecA**

*E. coli* RecA protein

**IPTG**

isopropyl-β-D(-)-thiogalactopyranoside

**ttRecA**

*T. thermophilus* RecA protein

**UV**

ultraviolet.

RecA protein is essential for homologous recombination and recombinational repair in *Escherichia coli* [1–3]. Many RecA homologs have been isolated not only from prokaryotes [4] but also from eukaryotes [5]. This suggests that the mechanism of DNA recombination is conserved in most living organisms, and that RecA protein may play a central role in the system. The properties of *E. coli* RecA protein have been extensively studied, and it has been shown that this relatively low-molecular-mass protein (38 kDa) has four major activities *in vitro* [1–3]. First, RecA monomers aggregate by themselves and polymerize on DNA to form a nucleoprotein helical filament. Second, RecA protein promotes a DNA strand exchange reaction. Third, RecA protein has single-stranded DNA (ssDNA)-dependent ATPase activity. Fourth, RecA protein has coprotease activity to induce an SOS response. The fact that these different activities are all shown by this single

Interestingly, poly(dA) and M13 ssDNA activated the ATPase at 65 °C but not at 37 °C. It has been reported that poly(dA) shows a positive band at around 192 nm by circular dichroism (CD) spectrometry [16]. This CD band in the region of vacuum UV indicates that poly(dA) does not have a random, but rather a specific, structure. In order to know the relationship between the structure and the ATPase activity, we measured the CD spectra of ssDNA and showed that poly(dA) and M13 ssDNA have a specific structure whereas poly(dC) and poly(dT) do not. We also showed that the positive CD band decreases as the temperature rises. From these observations, we surmise that the presence of M13 ssDNA prevents activation of ttRecA to express the ATPase activity.

Unlike the case of ttRecA, ecRecA expresses ATPase activity to varying degrees in the presence of various kinds of ssDNA. This observation suggests that the characteristic of ecRecA for ssDNA differs from that of ttRecA. The temperature dependency of the ATPase activity is similar to that of the structural alteration of poly(dA) and M13 ssDNA. In order to elucidate whether RecA protein can induce structural alteration of ssDNA, we carried out CD measurement of poly(dA) in the vacuum UV region in the presence of various concentrations of ecRecA at lower temperatures, and showed that ecRecA was able to change the specific structure of poly(dA) whereas ttRecA could not. From these observations, we discuss the difference between ecRecA and ttRecA with regard to the capability of modifying the specific structure of ssDNA necessary for the binding.

## Experimental procedures

### Enzymes and chemicals

The sources of enzymes and reagents were as follows: DNA modification enzymes including restriction enzymes were from Takara (Japan), Gene, Toyobo, Japan and New England Biolabs (USA); isopropyl- $\beta$ -D(-)-thiogalactopyranoside (IPTG) was from Wako Pure Chemical Industry Ltd. (Japan); phosphocellulose (type P11) and DEAE-cellulose (type DE52) were from Whatman Biochemicals (Germany); plastic-backed DEAE-cellulose sheets (MN-Polygram CEL300PEI/UV) were from Machery and Nagel (England); [ $\alpha$ -<sup>32</sup>P]ATP was from ICN (USA); protein kinase (type II) was from Sigma (USA); pig heart lactate dehydrogenase (grade II) was from Toyobo; adenosine 5'-O-(3-thiotriphosphate) was from Boehringer Mannheim (Germany); poly(dT), poly(dC), poly(dA) and poly(dA)·poly(dT) were from Pharmacia (USA); and M13 double-stranded DNA (dsDNA) were provided by T. Mikawa (Osaka University).

### Media, bacterial strains, plasmids and DNA manipulation

The *E. coli* strains used were DH5 $\alpha$  [18] for plasmid DNA preparation and BL21(DE3) harboring the pLysE plasmid [19] for expression of ttRecA. They were grown on Terrific broth or LB medium at 37 °C [18]. The plasmids used were pTA3, which contains the *recA* gene [9] in the pET3a expression vector [19]. DNA manipulation was carried out by standard procedures [18].

### Determination of nucleotide and polydeoxyribonucleic acid concentrations

The concentration of ATP or ATP- $\gamma$ -S was determined using the molar absorption coefficient of  $\epsilon_{259} = 15\,400\text{ M}^{-1}\cdot\text{cm}^{-1}$  or  $\epsilon_{264}$  respectively [20]. The residue molar concentrations of polydeoxyribonucleic acids were determined using the following molar absorption coefficients:  $\epsilon_{264} = 8520\text{ M}^{-1}\cdot\text{cm}^{-1}$  for poly(dT) [21];  $\epsilon_{268} = 7400\text{ M}^{-1}\cdot\text{cm}^{-1}$  for poly(dC) [21];  $\epsilon_{257} = 8600\text{ M}^{-1}\cdot\text{cm}^{-1}$  for poly(dA) [22];  $\epsilon_{264}$  for poly(dA)·poly(dT) [22].

### Overexpression of the *T. thermophilus recA* gene in *E. coli*

To create a *Nde* I restriction enzyme site in the first ATG codon of the *T. thermophilus recA* gene, a pair of DNA primers was synthesized (MilliGen/Biosearch, USA, Cyclone Plus DNA synthesizer). These were 5'-CCTGAGAGGTG CATATGGACGAGAG-3' and 5'-GAACTCCTTCTCA-3', the underlining indicating the *Nde* I site and the *Xmn* I site, respectively. A DNA fragment of about 8 kb was amplified by polymerase chain reaction under the same conditions as those described previously [9] using pTA3 as a template. The amplicon at the N-terminal side was cut with *Nde* I and *Xmn* I. The rest of the DNA fragment containing almost all of the *T. thermophilus recA* gene was prepared by cutting pTA3 with *Xmn* I and *Bam* HI. The pET3a vector was cut with *Nde* I and *Bam* HI. These three DNA fragments were ligated to construct the *T. thermophilus recA* gene under the regulation of the T7 promoter, and the resulting plasmid was named pTA3-*recA*. The sequence of the amplified region in pEA1 was confirmed by dideoxy sequencing using a 'Taq Dye Terminator Cycle Sequencing' kit (Applied Biosystems, USA).

## ATPase assay

At 37 °C, hydrolysis of ATP by ttRecA was measured by an enzyme coupling method [25, 26] at pH 7.5. A total of 200  $\mu\text{L}$  of contained 50 m M Tris/HCl, 10 m M  $\text{MgCl}_2$ , 100 m M KCl, 1 m M DTT, 10 m M phosphoenolpyruvate, 2 m M NADH, 25  $\text{U}\cdot\text{mL}^{-1}$  25 units $\cdot\text{mL}^{-1}$  lactate dehydrogenase, DNA, ATP and ttRecA as indicated. The activity was measured from the decrease in a 0.1-cm cell using a Hitachi spectrophotometer, model U-3000. Because the coupling enzymes are heat-inactivated at high temperature, a thin-layer chromatography (TLC) method [17] to measure ATPase activity at 65 °C. The procedure was described in the previous paper. Analysis of the ATPase activity was performed using the Hill equation (28):

where  $v$  and  $V_{\text{max}}$  are the initial and maximal initial ATPase reaction velocities,  $[S]$  is the concentration of substrate (ATP),  $K_d$  is the dissociation constant and  $n$  is the Hill coefficient. Eqn (1) was rearranged to give

Kinetic parameters were determined by linear fitting to Eqn (2) at various concentrations of ATP.

## Apparent dissociation constants between ttRecA and DNA

The concentration dependence of various kinds of DNA on the ATPase activity of ttRecA was measured. ATPase activity at 37 °C was measured by the enzyme coupling method with 5  $\mu\text{M}$  of ttRecA. The activity at 65 °C was measured by the TLC method with 2.5  $\mu\text{M}$  of ttRecA. To determine the intensity of binding between ttRecA and DNAs, we calculated the apparent dissociation constant from Eqn (3):

where E, L and EL are ttRecA, DNA, and their complex, respectively.  $K'$  is the dissociation constant defined by

where parentheses represent respective concentrations.

The total concentrations of ttRecA ( $[E]_0$ ) and DNA ( $[L]_0$ ) are expressed as follows:

The concentration of ttRecA-DNA complex is obtained from Eqns (4)–(6),


The apparent velocity of the reaction,  $v$ , is expressed as:

where  $k$  is a constant. The kinetic parameters were determined by fitting Eqn (8) to the data. A computer program, Igor (WaveMetrics) was used for calculation of the parameters.

## Strand exchange assay

A total of 20  $\mu\text{L}$  of reaction mixture contained 10 m M Tris/HCl (pH 7.5), 10 m M  $\text{MgCl}_2$ , 30 m M KCl, 20 m M ATP, 1 m M DTT, 10 m M ssDNA, 30  $\mu\text{M}$  M13mp18 dsDNA linearized by *Bam* HI digestion, and various concentrations of ttRecA. It also contained an enzyme coupling system (10 m M phosphoenolpyruvate, 25  $\text{U}\cdot\text{mL}^{-1}$  pyruvate kinase). The mixture was incubated at 37 °C or 65 °C for 30 min. The reaction was stopped by addition of a solution to the final concentration of 10 m M EDTA, 1% SDS and 0.2  $\text{mg}\cdot\text{mL}^{-1}$  proteinase K. A

mass of about 36 kDa, which agreed with that calculated from the amino acid sequence translated from its nucleotide sequence. The first 20 residues were M-D-E-S-K-R-K-A-L-E-N-A-L-K-A-I-E-K-E-F, which coincided with the sequence translated from the first ATG start codon of the *T. thermophilus recA* gene. To confirm the accurate molecular mass of ttRecA, the purified protein was analyzed by ion-spray mass spectrometry. The determined value, 36 385, agreed well with the calculated one, 36 384.



**Figure 1.**


[Open in figure viewer](#)

**Overproduction and purification of ttRecA.** The fractions obtained during the purification of ttRecA were electrophoresed on an SDS/polyacrylamide (12.5%) gel [46] and stained with Coomassie Brilliant Blue. Molecular mass markers (in kilodaltons) are shown on the left. Lane 1, total cell extract; lane 2, cleared lysate; lane 3, post-heat treatment at 65 °C; lane 4, post-phosphocellulose DEAE-cellulose. The arrow indicates the ttRecA. About 1–10 µg of protein was loaded into each lane of the gel.

## Kinetic parameters for ATPase activity of ttRecA

Including ecRecA, many proteins which have Walker's A-type nucleotide binding motif, GXXXXGK<sup>T</sup>/<sub>S</sub>[29], show ATPase activity with a conserved consensus sequence [9, 10] and ssDNA-dependent ATPase activity [10]. We studied the temperature dependence of ttRecA activity with poly(dT) as a ssDNA was measured using a TLC method. Its activity appeared maximal at 65 °C. At 37 °C, which is a temperature for many mesophilic organisms, the activity was about half that at the maximum. At over 90 °C and below 15 °C, no activity was observed. The optimum temperature of 65 °C for ttRecA activity is consistent with the fact that *T. thermophilus* can grow best at 65 °C [8].

Although some RecA proteins have been isolated from thermophiles [10, 30], no kinetic studies of their activities have been reported. We measured the ATPase activity of ttRecA at various concentrations of ATP to determine its kinetic parameters. For measurement of ATPase activity, the TLC method is advantageous when the reaction temperature is high, but disadvantageous for following the reaction in real time. On the other hand, the enzyme coupling method, which is now used widely for measuring ATPase activity, is useful for analysis of steady-state activity. In this study, the reaction can be followed spectroscopically in real time. However, the enzyme coupling method cannot be used at high temperatures because the coupling enzymes used here, lactate dehydrogenase (from pig heart) and pyruvate kinase (from rabbit muscle), are not thermophilic. Therefore, ATPase activity at 65 °C was measured using the TLC method. In the presence of poly(dT) as a ssDNA, the amount of ATP



**Figure 2.**

[Open in figure viewer](#)

**ATPase activities of ttRecA under various conditions.** (A) ATPase assays were performed by the TLC method at 65 °C as described in Experimental procedures. The reaction mixture contained 2.5 μM ttRecA and 100 μM poly(dT). (B) ATPase assays were performed by the enzyme coupling method at 37 °C as described in Experimental procedures. The reaction mixture contained 5 μM ttRecA and 100 μM poly(dT) (solid circles) or poly(dC) (clear circles). Substrate dissociation constant ( $K$ ), turnover number ( $k_{cat}$ ) and Hill coefficient ( $n$ ) for the hydrolysis of ATP were determined from the insets in the figures (see text for details).

**Table 1. Steady-state kinetic parameters for ATP hydrolysis catalyzed by RecA protein.**

	Temperature (°C)	ssDNA	$K^a$ (mM)	$k_{cat}$ (s <sup>-1</sup> )	$n^b$	$k_{cat}/K$ (s <sup>-1</sup> ·M <sup>-1</sup> )
ttRecA	65	poly(dT)	1.4	0.60	1.9	$4.3 \times 10^2$
	37	poly(dT)	0.34	0.28	2.1	$8.2 \times 10^2$
	37	poly(dC)	0.29	0.34	2.0	$1.2 \times 10^3$
ecRecA	37	poly(dT)	0.10	0.35	6.6	$3.5 \times 10^3$
	37	φX174 ssDNA	0.038	0.17	3.3	$4.5 \times 10^3$

<sup>a</sup>  $K$ , Substrate dissociation constant. <sup>b</sup>  $n$ , Hill coefficient. <sup>c</sup> All parameters for ttRecA protein were determined from [Fig. 2](#).

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.