



US006087099A

United States Patent [19]

Gupte et al.

[11] Patent Number: **6,087,099**

[45] Date of Patent: **Jul. 11, 2000**

[54] **METHOD FOR SEQUENCING BOTH STRANDS OF A DOUBLE STRANDED DNA IN A SINGLE SEQUENCING REACTION**

[75] Inventors: **Jamila Gupte**, Layton; **Arnold Oliphant**, Erda, both of Utah

[73] Assignee: **Myriad Genetics, Inc.**, Salt Lake City, Utah

[21] Appl. No.: **08/925,277**

[22] Filed: **Sep. 8, 1997**

[51] Int. Cl.⁷ **C12P 19/34; C07H 21/04**

[52] U.S. Cl. **435/6; 435/91.2; 536/24.3; 536/24.33; 536/26.6**

[58] Field of Search **536/24.33, 24.3, 536/26.6; 435/91.2, 6**

[56] **References Cited**

U.S. PATENT DOCUMENTS

5,411,875 5/1995 Jones 435/91.2

OTHER PUBLICATIONS

Chadwick, RB, Conrad, MP, McGinnis, MD, JohnstonDow, L, Spurgeon, SL and Kronick, MN (1996). "Heterozygote and Mutation Detection by Direct Automated Fluorescent DNA Sequencing Using a Mutant Taq DNA Polymerase", *BioTechniques* 20:676-683.

Choi, T-J, Wagner, JD and Jackson, AO (1994). "Sequence Analysis of the Trailer Region of Sonchus Yellow Net Virus Genomic RNA", *Virology* 202:33-40.

Jones, DH (1995). "Panhandle PCR", *PCR Methods and Applications* 4:S195-S201.

Jones, DH and Winistorfer, SC (1992). "Sequence specific generation of a DNA panhandle permits OCR amplification of unknown flanking DNA", *Nucleic Acids Research* 20:595-600.

Jones, DH and Winistorfer, SC (1993). "Genome Walking with 2-to 4-kb Steps Using Panhandle PCR", *PCR Methods and Applications* 2:197-203.

Ju, J., Ruan, C., Fuller, CW, Glazer, AN and Mathies, RA (1995). "Fluorescence energy transfer dye-labeled primers for DNA sequencing and analysis", *Proc. Natl. Acad. Sci. USA* 92:4347-4351.

Wetzel, T., Dietzgen, RG and Dale, JL (1994). "Genomic Organization of Lettuce Necrotic Yellows Rhabdovirus", *Virology* 200:401-412.

Primary Examiner—Ardin H. Marschel

Assistant Examiner—Joyce Tung

Attorney, Agent, or Firm—Rothwell, Figg, Ernst & Kurz PC

[57] **ABSTRACT**

A method is presented which uses a unique opposite strand joining strategy during PCR of an original DNA to generate a product which, when sequenced with a single sequencing primer yields the sequence of both strands of the original DNA. The PCR primers include 1) a modified oligomer corresponding to the 5' end of a first strand of the DNA to be amplified wherein said modified oligomer includes the reverse complementary sequence to a sequence within said first strand of DNA and a specific PCR priming sequence which will specifically hybridize to a portion of the DNA to be amplified and 2) a second oligomer corresponding to the 5' end of the second strand of the DNA to be amplified and which contains the priming sequence for the second strand of the DNA and will specifically hybridize to a portion of the DNA to be amplified. During PCR an intermediate product is formed where one end of one strand loops around to hybridize to its complement on the same strand. This results in a hairpin structure which elongates using its own strand as a template to form a double sized product that contains the sequence of both original strands. Upon denaturation this yields single strands with the single strands having the sequence of both of the original strands included in tandem. Sequencing these single strands using a single primer, e.g., a primer complementary to the second oligomer, yields the sequences of both strands of the DNA of interest.

6 Claims, 3 Drawing Sheets

19R,5' AGGAACAGCTATGACCATGATCCTCATATCATGGAATTTGT 3' SEQ ID NO:1
|-----M13R-----|-----G2'-----|

19P,5' GTTTCCAGTCAGCAGGTCATCTTCCTGCTCTTTTGT 3' SEQ ID NO:2
|-----M13P-----|-----G1-----|

19XP,5' CAGCGATTCCTCATCTTCCTGCTCTTTTGT 3' SEQ ID NO:3
|-----C'-----|-----G1-----|

5'...tcctctGTCATCTTCCTGCTCTTTTGTGAATCGCTaccctet...3 SEQ ID NO:4
|-----G1-----|-----C-----|

19R,5' **AGGAAACAGCTATGACCAT**TGATCCTCATTATCATGGAAAATTTGT 3' SEQ ID NO:1
|-----M13R'-----||-----G2'-----|

FIG. 1A

19F,5' **GTTTTCCCAGTCACGACG**GCATTCTTCCTGTGCTCTTTTGT 3' SEQ ID NO:2
|-----M13F'-----||-----G1'-----|


FIG. 1B

19XF,5' **CAGCGATT**CGTCATTCTTCCTGTGCTCTTTTGT 3' SEQ ID NO:3
|---C'---||-----G1'-----|

FIG. 1C

5' ...tctctGT**CATTCTTCCTGTGCTCTTTTGTGAATCGCTG**acctct...3' SEQ ID NO:4
|-----G1'-----||---C'---|

FIG. 1D






FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D

METHOD FOR SEQUENCING BOTH STRANDS OF A DOUBLE STRANDED DNA IN A SINGLE SEQUENCING REACTION

BACKGROUND OF THE INVENTION

Sequencing of nucleic acids is an extremely important and widely used technique. It is used for a variety of purposes. One such purpose is to identify whether mutations within genes of known sequence are present in a sample of DNA taken from a person. This is especially important in diagnosing whether the person may have a disease which is known to be associated with specific mutations in the gene being analyzed. When this type of testing is performed, it is common to sequence both strands of DNA to minimize any errors which may occur in the sequencing. To date, when sequencing both strands by the Sanger dideoxy method there has been a requirement to use one primer to sequence the sense strand and a second primer to sequence the antisense strand of the double-stranded DNA. The two strands have been sequenced in separate sets of reactions. The present invention is a technique by which both strands of DNA are sequenced in a single set of reactions using only a single primer. This method allows one to use fewer reactions for obtaining the data. This is especially important for laboratories which will be processing many samples. The use of fewer reactions will decrease the cost of analysis.

DNA sequencing methods were developed during the 1970s by Maxam and Gilbert (1977) and by Sanger (1977). The Sanger method which uses dideoxy nucleotides to terminate newly synthesized DNA strands is most commonly used and has been adapted such that it can be used with fluorescent markers rather than radioactivity. One variation is a technique called cycle sequencing in which DNA sequencing is combined with polymerase chain reaction (PCR). Chadwick et al. (1996) teach a variation of cycle sequencing in which a mutant Taq DNA polymerase is utilized.

The polymerase chain reaction itself is only one of a number of different methods now available for amplifying nucleic acids. Some of the other methods include ligase chain reaction (Wu and Wallace, 1989), Strand Displacement Amplification (SDA) (Walker, U.S. Pat. No. 5,455,166 (1995); Walker et al., 1992), thermophilic SDA (Spargo et al., 1996), and 3SR or NASBA (Compton, 1991; Fahy et al., 1991).

The instant invention is a method of using a specially designed oligomer which contains a reverse complement sequence along with a standard primer during PCR. This generates a double stranded DNA product such that when it is denatured one end of the resulting single stranded DNA loops around to form an intrastrand stem-loop structure. This structure is then elongated thereby producing a double-stranded DNA but wherein the two strands are joined by a loop. This method is referred to as opposite strand joining PCR. When denatured this product forms a single-stranded DNA which contains both strands of the original DNA. When this resulting single-stranded DNA is sequenced it yields the sequence of both strands of the original double-stranded DNA.

A similar stem-loop DNA structure was used as a template for PCR amplification by Jones et al. (1992). The Jones et al. reference describes a "panhandle PCR" method. This technique introduced a self-complementary portion into the target DNA strand by ligation. The goal of panhandle PCR is to amplify unknown sequence by generating a stem loop template structure for PCR whereas one of the goals of

opposite strand joining PCR is to amplify known sequence by generating a stem-loop structure during PCR and then sequencing both strands of the longer product in one sequencing reaction. Another use for opposite strand joining PCR is in denaturing gradient gel electrophoresis techniques wherein the use of this technique can form a covalently bonded hairpin loop which can replace the use of a GC clamp. Yet another use for opposite strand joining PCR is simply the use of the method effectively to join together the two strands of any double stranded DNA into a single strand of DNA for any desired purpose.

SUMMARY OF THE INVENTION

A method is presented which uses a unique opposite strand joining strategy during PCR of an original DNA to generate a product which, when sequenced with a single sequencing primer yields the sequence of both strands of the original DNA. The PCR primers include 1) a modified oligomer corresponding to the 5' end of a first strand of the DNA to be amplified wherein said modified oligomer includes the reverse complementary sequence to a sequence within said first strand of DNA and a specific PCR priming sequence which will specifically hybridize to a portion of the DNA to be amplified and 2) a second oligomer corresponding to the 5' end of the second strand of the DNA to be amplified and which contains the priming sequence for the second strand of the DNA and will specifically hybridize to a portion of the DNA to be amplified. During PCR an intermediate product is formed where one end of one strand loops around to hybridize to its complement on the same strand. This results in a hairpin structure which elongates using its own strand as a template to form a double sized product that contains the sequence of both original strands. Upon denaturation this yields a single strand having the sequence of both of the original strands included in tandem. Sequencing these single strands using a single primer, e.g., a primer complementary to the second oligomer, yields the sequences of both strands of the DNA of interest.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A-1D show the primer design used in the Example. FIG. 1A shows the sequence of primer 19R which consists of the -28M13 reverse DET primer sequence (shown in bold) which is 5' to the gene specific sequence G2'. FIG. 1B shows the sequence of primer 19F which consists of the -40M13 forward DET sequence (shown in bold) which is 5' to the gene specific sequence G1. FIG. 1C shows the sequence of the opposite strand joining primer 19XF which consists of a short reverse complemented genomic sequence C' (shown in bold) which is 5' to the gene specific sequence G1 used in primer 19F. FIG. 1D shows the genomic sequence in the region of the opposite strand joining primer. The gene specific sequence G1 (shown in nonbolded upper case letters) used in both the 19F and 19XF primers is 5' of sequence C (shown in bold upper case letters). It is this genomic region C which is reverse complemented (and therefore called C') and placed 5' to the gene specific sequence G1 in the opposite strand joining primer 19XF.

FIGS. 2A-2F illustrate the opposite strand joining strategy. Throughout these figures, all the strands labeled •SEQ are substrates for dye primer sequencing.

FIG. 2A shows genomic DNA in the region of exon 19. This is shown as four sections on each strand with one strand having G1, C, the exon 19 containing region, and G2 and the opposite strand being designated with primes, e.g., G1', C',

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.