

PHARMACEUTICAL PREFORMULATION: The Physicochemical Properties of Drug Substances

JAMES I. WELLS, B.Sc., M.Sc., Ph.D., M.P.S., MInstPkg.
Development Group Leader
Pharmaceutical Research and Development
Pfizer Central Research, Sandwich, Kent



ELLIS HORWOOD LIMITED
Publishers • Chichester

Halsted Press: a division of
JOHN WILEY & SONS

QV
744
W454p
1988

First published in 1988 by

ELLIS HORWOOD LIMITED

Market Cross House, Cooper Street,
Chichester, West Sussex, PO19 1EB, England

The publisher's colophon is reproduced from James Gillison's drawing of the ancient Market Cross, Chichester.

Distributors:

Australia and New Zealand:

JACARANDA WILEY LIMITED

GPO Box 859, Brisbane, Queensland 4001, Australia

Canada:

JOHN WILEY & SONS CANADA LIMITED

22 Worcester Road, Rexdale, Ontario, Canada

Europe and Africa:

JOHN WILEY & SONS LIMITED

Baffins Lane, Chichester, West Sussex, England

North and South America and the rest of the world:

Halsted Press: a division of

JOHN WILEY & SONS

605 Third Avenue, New York, NY 10158, USA

South-East Asia

JOHN WILEY & SONS (SEA) PTE LIMITED

37 Jalan Pemimpin # 05-04

Block B, Union Industrial Building, Singapore 2057

Indian Subcontinent

WILEY EASTERN LIMITED

4835/24 Ansari Road

Daryaganj, New Delhi 110002, India

© 1988 J. I. Wells/Ellis Horwood Limited

British Library Cataloguing in Publication Data

Wells, James I. (James Ingram), 1950-

Pharmaceutical preformulation.

1. Drugs. Physiochemical aspects .

I. Title

615'.19

Library of Congress Card No. 88-9233

ISBN 0-7458-0276-1 (Ellis Horwood Limited)

ISBN 0-470-21114-8 (Halsted Press)

Phototypeset in Times by Ellis Horwood Limited

Printed in Great Britain by Hartnolls, Bodmin

COPYRIGHT NOTICE

All Rights Reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the permission of Ellis Horwood Limited, Market Cross House, Cooper Street, Chichester, West

2

Solubility

When a preformulation programme begins, the availability of bulk is always limited and the scientist may only have 50 mg. Thus, it is imperative that the best use of the limited bulk is made, to support the continuing efforts of the synthetic chemists and the biologists, pursuing activity and toxicity screens. Furthermore, because the compound is new, the quality is invariably poor, so that a number of impurities may be present and often the first seeding comes down as a metastable polymorph (Chapter 3). Accordingly if nothing else is measured, the solubility and pK_a must be determined since these largely control all future work. The solubility dictates the ease with which formulations for gavage and intravenous injection studies in animals are obtained. The pK_a allows the informed use of pH to manipulate solubility and choose salts, should they be required to achieve good bioavailability from the solid state, and to improve stability (Chapter 5) and powder properties (Chapter 6).

Kaplan (1972) suggested that, unless a compound has an aqueous solubility in excess of 1% (10 mg ml^{-1}) over the pH range 1–7 at 37°C , then potential adsorption problems may occur. He also found that if the intrinsic dissolution rate (IDR, section 2.7) was greater than $1 \text{ mg cm}^{-2} \text{ min}^{-1}$ then adsorption was unimpeded, while less than $0.1 \text{ mg cm}^{-2} \text{ min}^{-1}$ gave dissolution rate-limited adsorption. This ten-fold difference in dissolution rate translates to a lower limit for solubility of 1 mg ml^{-1} since under sink conditions, dissolution rate and solubility are proportional (Hamlin *et al.*, 1965). A solubility of less than 1 mg ml^{-1} indicates the need for a salt, particularly if the drug is to be formulated as a tablet or capsule. In the range 1–10 mg ml^{-1} , serious consideration should be given to salt preparation. These guidelines are shown graphically in Fig. 2.1. Where the solubility of the drug cannot be manipulated in this way (a neutral molecule: glycoside, steroid, alcohol or where the pK_a is less than 3 for a base or greater than 10 for an acid), then liquid filling (a solution in PEG 400, glyceryl triacetate or fractionated coconut oil) in a soft gelatin capsule, or as a paste or semisolid (dissolved in oil or triglyceride) in a hard gelatin capsule may be necessary.

2.1 INTRINSIC SOLUBILITY (C_0)

First examine the chemical structure, or determine the solubility in 0.1 N HCl, 0.1 N NaOH and water by UV. An increase in acid over aqueous solubility suggests a weak




Fig. 2.1 — Relationship between drug pK_a and solubility. Solubility $> 10 \text{ mg ml}^{-1}$ at $< \text{pH } 7$ (Kaplan, 1972) is required to ensure good bioavailability.

base, and in alkali, a weak acid. In both cases, a dissociation constant (pK_a) will be measurable and salts should form. An increase in both acid and alkali solubility suggests either amphoteric or zwitterionic behaviour and there will be at least two pK_a s, one acidic and one basic. No change in solubility suggests a non-ionizable,

neutral molecule with no measurable pK_a . Here solubility manipulation will require either solvents or complexation.

When the purity of the drug sample can be assured, then the value obtained in acid for a weak acid or alkali for a weak base can be assumed to be the intrinsic solubility (C_o), i.e. the unionized form. However, since absolute purity is often in doubt on the first few synthetic batches, it is more accurate to determine this crucial solubility from a phase-solubility diagram (Fig. 2.2). The solubility should ideally be measured at two temperatures:

- (1) 4°C: to ensure good physical stability and extended short-term storage and chemical stability until more definitive data are available. The density of water is maximum at 4°C and this imposes the greatest challenge to saturated aqueous solubility.
- (2) 37°C: to support biopharmaceutical evaluation, since this is body temperature.




Fig. 2.2 — Effect of drug : solvent ratio on solubility when the drug is impure.

Assuming the compound is a base and the estimate in 0.1 N NaOH gave 1 mg ml^{-1} , then four solutions of 3 ml should be set up containing 3, 6, 12 and 24 mg of drug respectively. These give the phase ratios shown in Fig. 2.1. Three millilitres is the smallest volume which can be manipulated, by either centrifugation or filtration, followed by dilution for UV analysis. The vials containing the samples should be agitated continuously for 16 hours (overnight) and then the concentration in solution determined. The data should be plotted according to Fig. 2.1 and the line extrapolated to zero phase ratio, where the 'solubility' will be independent of solvent level

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.