

Neuromuscular Electrical Stimulation for Quadriceps Muscle Strengthening After Bilateral Total Knee Arthroplasty: A Case Series

Jennifer E. Stevens, PT, PhD¹

Ryan L. Mizner, MPT²

Lynn Snyder-Mackler, PT, ScD, ATC, SCS, FAPTA³

Study Design: A case series.

Objectives: The purpose of this case series was to assess the effect of high-intensity neuromuscular electrical stimulation (NMES) on quadriceps strength and voluntary activation following total knee arthroplasty (TKA).

Background: Following TKA, patients exhibit long-term weakness of the quadriceps and diminished functional capacity compared to age-matched healthy controls. The pain and swelling that results from surgery may contribute to quadriceps weakness. The use of high-intensity NMES has previously been shown to be effective in quickly restoring quadriceps strength in patients with weakness after surgery.

Methods and Measures: All patients were treated for 6 weeks, 2 to 3 visits per week, in outpatient rehabilitation. Five patients (NMES group) participated in a voluntary exercise program for both knees and NMES for the weaker knee. Three patients (exercise group) participated in a voluntary exercise program for both knees without NMES. For each treatment session, 10 isometric electrically elicited muscle contractions were administered at maximally tolerated doses to the initially weaker leg of the NMES group. Quadriceps strength and muscle activation were repeatedly assessed up to 6 months after surgery using burst superimposition techniques.

Results: At 6 months, the weak NMES-treated legs of 4 of 5 patients in the NMES group had surpassed the strength of the contralateral leg. In contrast, none of the weak legs in the exercise group were stronger than the contralateral leg at 6 months. Changes in quadriceps muscle activation mirrored the changes exhibited in strength.

Conclusion: When NMES was added to a voluntary exercise program, deficits in quadriceps muscle strength and activation resolved quickly after TKA. *J Orthop Sports Phys Ther* 2004;34:21-29.

Key Words: geriatric, inhibition, rehabilitation, total knee replacement

Simultaneous bilateral total knee arthroplasty (TKA) has become an increasingly common surgical procedure for patients with bilateral knee osteoarthritis or rheumatoid arthritis.^{1,5,9,20,21,23} Typically, patients are only permitted to undergo simultaneous bilateral TKA if they are in excellent health due to the increased risk for complications from 2 surgical procedures (eg, longer anesthesia times).^{1,20,23} When patients are appropriately chosen for the procedure, outcomes are comparable to patients who undergo a staged bilateral knee replacement,²³ yet the total rehabilitation time is less.²⁰

Patients after TKA are often plagued with quadriceps femoris weakness and functional deficits that continue for years after surgery.^{2,4,15,39} These limitations in strength and function are thought to result from a combination of preexisting weakness before surgery and pain and swelling postsurgery.^{18,22,26} After bilateral TKA, patients are doubly affected by this persistent quadriceps weakness, which may result in serious functional limitations. Quadriceps weakness has been correlated with

¹ Postdoctoral Associate, Department of Physical Therapy, University of Florida, Gainesville, FL. Dr. Stevens was a doctoral student, Biomechanics and Movement Science Program, University of Delaware, Newark, DE at the time of this study.

² Doctoral student, Biomechanics and Movement Science Program, University of Delaware, Newark, DE.

³ Professor, Department of Physical Therapy and Biomechanics and Movement Science Program, University of Delaware, Newark, DE.

This study was supported by NIH grant 1R01HD041055-01A1, NIH training grant T32 HD07490, and APTA Foundation Scholarships to Jennifer Stevens and Ryan Mizner (PODS). Research protocol was approved by the Human Subjects Review Committee at the University of Delaware.

Please address all correspondence to Lynn Snyder-Mackler, Department of Physical Therapy, 301 McKinly Laboratory, University of Delaware, Newark, DE 19716. E-mail: smack@udel.edu

and increased risk of falls.¹⁰ One source of this weakness is voluntary activation deficits, which result from a failure of recruiting all available motor units or a reduction in the maximal motor unit discharge rate from those motor units which are recruited.¹⁹ Voluntary activation deficit can be quantified by superimposing a supramaximal electrical stimulus on a maximally contracting muscle.^{10,19,34,36,37,41} Using this technique, voluntary activation deficits have been reported in patients with osteoarthritis or following knee surgery, including TKA.^{18,25,27,28,35} Voluntary activation deficits can result from pain,²⁷ effusion,¹² and joint damage,¹⁶ all of which are potentially present in patients after TKA. Voluntary exercise programs have had limited success in restoring quadriceps strength when a substantial voluntary activation deficit is present.¹⁷

Studies of various younger adult populations have demonstrated that neuromuscular electrical stimulation (NMES), at sufficient intensities, can be combined with volitional exercise to more effectively increase muscle strength and functional performance than volitional exercise alone.^{11,31,33} A recent case report suggested that incorporating high-intensity NMES into an outpatient rehabilitation program for muscle strengthening offers promise for restoring quadriceps strength.²¹ NMES has the potential to override muscle activation deficits resulting from impairments in central nervous system processing. In addition, NMES activates a greater proportion of type II (fast-twitch) muscle fibers when compared to volitional exercise at a comparable intensity.^{3,7,13,30,38} Type II fibers are essential for higher levels of force production and their activation may translate to improved functional performance.

Applying high-intensity electrical stimulation to only 1 limb following bilateral TKA provides an ideal opportunity for investigating the efficacy of NMES, as each patient provides a control limb for comparison. The purpose of this case series was to assess the effect of adding high-intensity NMES to a volitional strengthening program following TKA. The hypothesis was that adding NMES to a 6-week, voluntary exercise program would be more effective in improv-

METHODS

Patients

Eight patients with simultaneous, primary bilateral total knee replacements (Table 1) were recruited from local surgeons who performed tricompartmental cemented TKA with a medial parapatellar approach. Patients were excluded from the study if they were diagnosed with diabetes mellitus, uncontrolled blood pressure, neurological disorders, neoplasms, or had a body mass index (BMI = weight [kg]/height [m²]) of greater than 40 (morbidly obese).

Patients were assigned to 1 of 2 intervention groups: (1) "Ex legs" or (2) "NMES legs." Patients in the Ex legs group received the same intervention for both legs, which consisted of voluntary exercises for strengthening of the lower extremity. Patients in the NMES legs group participated in a voluntary exercise program for the leg that was stronger at initial evaluation, and NMES was added to the voluntary exercise program for the leg that was weaker at initial evaluation (Table 2). The Human Subjects Review Committee at the University of Delaware approved the study and all subjects gave informed consent.

Intervention

Intervention began 3 to 4 weeks following TKA, after staples were removed. All patients were scheduled for treatment at the University of Delaware Physical Therapy Clinic 3 times a week for 6 weeks (total of 18 visits) and were allowed to miss up to 4 visits. All in-clinic and home exercise interventions were documented in patient data booklets to ensure consistency of care and patient compliance. Rehabilitation for all patients included interventions to reduce pain and inflammation, improve incision mobility, restore knee flexion and extension range of motion, improve bilateral lower-extremity strength

TABLE 1. Patient demographics for patients with bilateral total knee arthroplasty.

Patient	Height (m)	Weight (kg)	BMI	Age (y)	Gender	Group
1	1.75	103.4	33.8	64	M	NMES
2	1.78	97.1	30.8	67	M	NMES
3	1.85	93.0	27.1	75	M	NMES
4	1.68	88.9	31.5	64	M	NMES
5	1.85	99.8	29.2	63	M	NMES
6	1.73	118.8	39.7	76	F	Exercise
7	1.73	93.0	31.1	69	M	Exercise
8	1.78	87.5	27.8	61	M	Exercise

Abbreviations: BMI, body mass index (weight [kg]/height [m²]); F, female; M, male; NMES, neuromuscular electrical stimulation.

Group Name/Leg	Intervention
NMES group	
Weak NMES legs	Volitional exercise, NMES
Strong NMES legs	Volitional exercise
Exercise group	
Weak Ex legs	Volitional exercise
Strong Ex legs	Volitional exercise

Abbreviation: NMES, neuromuscular electrical stimulation.

using high-intensity voluntary exercises, and improve functional performance (see Table 3). Each patient was evaluated and treated on the basis of individual impairments, as the guidelines for intervention in Table 3 permitted.

NMES The weak NMES legs received identical intervention to all other legs except that 10 NMES-elicited quadriceps contractions were added to each treatment session. For the NMES intervention, patients were seated on an electromechanical dynamometer (KinCom; Chattanooga Corporation, Chattanooga, TN), and a measuring arm, which restrained the movement of the leg, was secured to the lower leg (Figure 1). The knee was positioned at 60° of knee flexion and all contractions were isometric. The patient's maximal voluntary isometric contraction (MVIC) was determined before the NMES intervention using the average peak force of two 5-second isometric contractions.

Self-adherent, flexible electrodes (7.6×12.7 cm) were placed over the subject's quadriceps to apply the electrical stimulation.²⁹ A clinical neuromuscular electrical stimulator (Versastim 380; Electromed Health Industries, Miami Beach, FL) was set to deliver a 2500-Hz alternating current, modulated at 50 bursts per second, with a ramp-up time of 2 to 3 seconds.²⁹ The intensity was set to the maximum intensity tolerated by the patient during each session. The patient was instructed to relax during stimulation and prevent cocontraction of the hamstrings as well as inadvertent voluntary quadriceps muscle contraction. The intensity was increased as tolerated throughout each session. Ten 10-second isometric contractions were elicited with an 80-second rest between each contraction.²⁹ The electrically elicited knee extension peak force produced by each contraction of the quadriceps muscle was recorded by customized software (LabVIEW V 4.0.1; National Instruments, Austin, TX) on a personal computer.

Testing Procedure

Quadriceps strength and voluntary muscle activation were tested. Testers were isolated from the subjects' interventions and were not aware of which patients received the NMES intervention. Testing was performed in the 3rd (initial evaluation), 6th

follow-up), and 24th (6-month follow-up) weeks after TKA. To minimize apprehension and discomfort during testing, patients were allowed to choose which leg was tested first during the initial evaluation. The order of testing remained the same for each subsequent testing session.

Quadriceps Strength and Activation Testing

Patients were seated on the electromechanical dynamometer with their hips flexed to approximately 85° and their knees flexed to 75°. The axis of the dynamometer was aligned with the axis of rotation of the knee joint and the bottom of the force transducer pad was positioned against the anterior aspect of the leg proximal to the lateral malleolus. The lower leg, thigh, and pelvis were stabilized using inelastic straps with Velcro closures. Two 7.6×12.7-cm self-adhesive neuromuscular stimulation electrodes were secured to the thigh. The anode was placed over the motor point of the rectus femoris and the cathode was placed over the motor point of the vastus medialis.

Strength testing began by instructing all patients to perform three 3- to 5-second voluntary isometric contractions at an intensity that they perceived as 50% to 75% of their maximal effort. These contractions served to familiarize the patient with the apparatus and to warm up the muscle. In addition, the quadriceps femoris muscle was stimulated several times at intensities that would elicit muscle contractions to familiarize the patient with the sensation of electrical stimulation. After the patient was familiar with the procedure and the muscle warmed up, testing began by having the patient attempt a 3- to



FIGURE 1. Experimental setup for neuromuscular electrical stimulation (NMES) treatment. Patients were seated and stabilized with the hip flexed to 85° and the knee flexed to 75°. Two 7.62×12.70-cm electrodes were placed over the vastus medialis and proximal rectus femoris of the quadriceps.

Range of motion (ROM)	<ul style="list-style-type: none"> • Exercise bike (10-15 min); started with forward and backward pedaling with no resistance until enough ROM for full revolution; progression: lower seat height to produce a stretch with each revolution. • Active-assistive ROM for knee flexion, sitting or supine, using other leg to assist. • Knee extension stretch with manual pressure (in clinic) or weights (at home). • Patellar mobilizations: 3×30 superior/inferior; medial/lateral, as necessary.
Strength	<ul style="list-style-type: none"> • Quad sets, straight leg raises (without quad lag), hip abduction (sidelying), standing hamstring curls, seated knee extension, standing terminal knee extensions from 45° to 0°, step-ups (5.08-15.24-cm block), wall slides to 45° knee flexion; 1 to 3 sets of 10 repetitions for all strengthening exercises. • Criteria for progression: exercises are to be progressed (eg, weights, step height, etc) once the patient can complete the exercise correctly and feels maximally fatigued at the end of each set. • Progression: 0.454-0.907-kg weights added to exercises, step-downs (5.08-15.24-cm block), front lunges, wall slides towards 90° knee flexion.
Pain and swelling Incision mobility Functional activities	<ul style="list-style-type: none"> • Ice and compression as needed. • Soft tissue mobilization until incision moves freely over subcutaneous tissue. • Ambulation training with assistive device as appropriate with emphasis on heel strike, push-off at toe-off and normal knee joint excursions. • Emphasis on heel strike, push-off at toe-off and normal knee joint excursions when able to walk without assistive device. • Stair ascending and descending step over step when patient has sufficient concentric/eccentric strength.
Monitoring vital signs	<ul style="list-style-type: none"> • Blood pressure and heart rate are monitored at initial evaluation and as appropriate.

5-second MVIC while receiving verbal encouragement from the tester and visual feedback from the dynamometer's real time force display. During the contraction, a 135-V, 10-pulse, 100-pps train (1000-microsecond pulse duration) was delivered to the muscle to assess whether the subject was indeed maximally activating the quadriceps muscle (burst superimposition technique).^{19,24,34,37} A S8800 stimulator with a SIU8T stimulus isolation unit (Grass Instruments, Inc., Quincy, MA) was driven by a personal computer that uses customized software (LabVIEW V 4.0.1) to control the timing parameters of the stimulation protocol. Data were digitized at 200 Hz and analyzed with customized software (LabVIEW V 4.0.1). With full voluntary muscle activation, no increase in force was measured. If a subject was unable to activate the quadriceps muscle fully, MVIC testing was repeated up to 2 additional times and the maximal force noted during these 3 tests was used for analysis. Each attempt at achieving an MVIC was separated by 5 minutes to minimize the effects of muscle fatigue. The greatest maximal voluntary effort achieved during the testing session for each leg was used for analysis. The burst superimposition technique has been shown to be highly reliable in subjects without pathology, with repeated testing that demonstrated an intraclass correlation coefficient of 0.98 (mean age, 24.2 years; age range, 17-32 years).³²

Data Management and Analysis

Because of the limited number of subjects included in this study, the data are presented individually and qualitatively to describe trends in outcomes.

Quadriceps Strength The force (N) of the quadriceps MVIC was normalized to body mass index (N/BMI) for all comparisons to account for variations in force production that are a result of differences in body size. Three sets of comparisons were made for the 8 individuals: (1) comparison within the NMES legs and Ex legs groups (strong versus weak legs); (2) comparison of weak legs between groups (weak NMES legs versus weak Ex legs); (3) comparison of strong legs between groups (strong NMES legs versus strong Ex legs).

The average percent increase in normalized quadriceps muscle strength was also used to describe strength gains of the individuals. The percent increase was calculated using the following relationship:

$$\text{Percent increase in strength} = \frac{\text{6-month MVIC (N)} - \text{initial MVIC (N)}}{\text{initial MVIC (N)}} \times 100$$

Quadriceps Voluntary Muscle Activation Voluntary quadriceps muscle activation was calculated by measuring the peak volitional and electrically elicited forces during the MVIC test. The central activation ratio (CAR) was used to quantify voluntary muscle activation.¹⁹ The CAR is calculated by dividing the maximum voluntary force produced before the electrical stimulus by the maximum force produced during the superimposition of the electrical stimulus. A CAR of 1.0 signifies complete activation. In contrast, a CAR of less than 1.0 suggests incomplete voluntary muscle activation. The same comparisons outlined earlier for quadriceps strength were made for voluntary activation.

ation was also used to describe improvements in activation of the individuals. The percent increase was calculated using the following relationship:

$$\text{Percent increase in CAR} = \frac{6\text{-month CAR} - \text{initial CAR}}{\text{initial CAR}} \times 100$$

NMES Dose The daily NMES dose was calculated as a ratio of the highest electrically elicited knee extension force obtained in a session by the maximal MVIC achieved on that same day. The NMES doses are reported as average doses obtained over the 6 weeks of treatment.

RESULTS

Compliance

The average number of sessions was 17. No patient missed more than 3 sessions.

Quadriceps Strength and Voluntary Muscle Activation

Strong Versus Weak Legs The percent increases in quadriceps strength for the weak NMES legs ranged from 221% to 451% as compared to 50% to 152% for the strong NMES legs; whereas the percent increases in quadriceps strength for the weak Ex legs ranged from 41% to 148% as compared to 30% to 71% for the strong Ex legs (Table 4). Initially, the average quadriceps strength of almost all legs receiving NMES was less than that of the legs that received only voluntary exercise for strengthening (Table 5).

After 3 weeks of treatment (midtraining), the quadriceps strength of the weak NMES legs showed a dramatic improvement in 4 out of 5 patients and was almost identical to, or had surpassed, the strength of the contralateral strong NMES legs. At 6 months, 4 out of 5 patients with initially weak NMES legs remained stronger than their initially stronger

weak Ex legs showed a similar initial jump in quadriceps strength within the first 3 weeks of training (patient 6), and none of the weak Ex legs were stronger as compared to their contralateral leg at 6 months (Table 5).

Quadriceps strength testing demonstrated consistent linear or curvilinear increases in strength for all legs over the course of 6 months, with the exception of patient 6 (Table 5). Patient 6 showed the greatest amount of variability in force production during consecutive testing sessions and had greater oscillations in strength than the other patients.

Quadriceps voluntary activation for the legs receiving stimulation underwent a concurrent increase compared to the legs that did not undergo NMES during the first 3 weeks of treatment (Table 6). Quadriceps voluntary activation did not change as dramatically during the remaining testing sessions as it did during the first 3 weeks of training for patients in the NMES group.

Weak NMES Legs Versus Weak Ex Legs Initial quadriceps strength of the weak NMES legs was less for patients 1 through 5 than that of the weak Ex legs of patients 7 and 8 (Table 5). Patient 6 was the exception to this observation and had quadriceps weakness that was comparable to that of the weak NMES legs. By midtraining, quadriceps strength of the weak NMES legs had almost equaled or surpassed that of the weak Ex legs in all patients except patient 3. Quadriceps strength of the weak Ex legs improved by midtraining. Both the weak NMES legs and weak Ex legs continued to show improvements in strength through the 6-month follow-up, but the weak NMES legs demonstrated the greatest overall improvement (Table 5).

Voluntary muscle activation of the quadriceps increased for all the weak NMES legs within the first 3 weeks of treatment (Table 6). In comparison, the weak Ex legs only had 1 substantial change in quadriceps voluntary activation in patient 8, which

TABLE 4. Percent change in quadriceps strength and voluntary activation from initial evaluation to 6-month follow-up.

Patient	Increase in Quadriceps Strength (%)		Increase in CAR (%)	
	Weak Legs	Strong Legs	Weak Legs	Strong Legs
1*	451	76	39.0	-1.6
2*	302	121	87.0	11.0
3*	233	152	66.0	52.0
4*	351	133	34.0	5.0
5*	221	50	34.0	-3.0
6†	41	30	-6.0	-14.0
7†	80	67	-0.8	-4.0
8†	148	71	54.0	3.0

Abbreviation: CAR, central activation ratio.

*Neuromuscular electrical stimulation group.

†Exercise group.



Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.