

## Short Communication

# KINETIC ISOTOPE EFFECTS IMPLICATE A SINGLE OXIDANT FOR CYTOCHROME P450-MEDIATED O-DEALKYLATION, N-OXYGENATION, AND AROMATIC HYDROXYLATION OF 6-METHOXYQUINOLINE

Received March 23, 2006; accepted May 12, 2006

### ABSTRACT:

One major point of controversy in the area of cytochrome P450 (P450)-mediated oxidation reactions is the nature of the active-oxygen species. A number of hypotheses have been advanced which implicate a second oxidant besides the iron-oxo species designated as compound I (Cpd 1). This oxygen is thought to be either an iron-hydroperoxy species (Cpd 0) or a second spin-state of Cpd 1. Very little information is available on what fraction of P450 oxidations is mediated by the two different oxidants. Herein, we report results on three cytochrome P450-mediated reactions: O-dealkylation, N-oxygenation, and aromatic hydroxylation, which occur by three distinct chemical mechanisms. We have used ki-

netic isotope effects to test for branching from O-demethylation to N-oxygenation and aromatic hydroxylation, using 6-methoxyquinoline and  $^2\text{H}_3$ -6-methoxyquinoline as substrates for P4501A2. Identical large inverse isotope effects on  $V_{\text{max}}/K_m$  are obtained for the formation of both the N-oxide and the phenol. This indicates that all three reactions occur through the same enzyme-substrate complex and, thus, through a single iron-oxygen species. The nature of the iron-oxygen species is less certain but is more likely to be iron-oxo Cpd 1, given the energetics of these reactions.

The ubiquitous cytochrome P450 enzymes (P450s) are a superfamily of monooxygenases that function to metabolize a variety of endogenous and xenobiotic compounds (Ortiz de Montellano, 1995; Meunier et al., 2004). A great deal of effort has been expended on probing the oxygenation mechanisms and identifying the intermediate species formed in the catalytic cycle of the enzyme. Until recently, a consensus seemed to have been reached that the high-valent iron-oxo species (Cpd 1) is the sole active oxygenating species in P450 (Groves et al., 1978; Guengerich and MacDonald, 1990; Ortiz de Montellano, 1995). This mechanism was first elaborated by Groves et al. (1978) for aliphatic substrates as being an initial hydrogen atom abstraction followed by recombination to produce an alcohol. However, many recent publications implicate an iron-hydroperoxy species (Cpd 0) as a second electrophilic oxidant (Vaz et al., 1998; Hutzler et al., 2003; Chandrasena et al., 2004). The hypothesis that Cpd 0 is a second oxidant in the P450 catalytic cycle began with a series of site-directed mutagenesis experiments performed by Coon and coworkers (Vaz et al., 1998) in which an active-site threonine is replaced by an alanine. The mutation is thought to affect proton delivery that converts Cpd 0 to Cpd 1. The results indicate that some of the mutant enzymes will perform olefin epoxidation at enhanced rates while decreasing aliphatic C-H hydroxylation (Vaz et al., 1998). Radical probe and kinetic isotope effect experiments were used by Newcomb and coworkers (Chandrasena et al., 2004) to conclude that Cpd 0 is the predominant

oxidant affecting aliphatic hydroxylation of substituted cyclopropanes but that Cpd 1 oxygenates higher barrier reactions such as hydroxylation of the methyl group of straight-chain aliphatic compounds. In separate studies, Hutzler et al. (2003) have suggested that two different oxidants may be involved in O-dealkylation and N-dealkylation. Jin et al. (2003) came to the conclusion that Cpd 0 could epoxidize alkenes but not hydroxylate the high-energy aliphatic C-H bond on camphor, and Volz et al. (2002) suggested that Cpd 0 and 1 may be the sole oxidants involved in N-dealkylation and sulfoxidation, respectively.

An alternative explanation for these results, as proposed by Shaik and coworkers (Ogliaro et al., 2002), is a two-state reactivity model in which Cpd 1 has two accessible spin states that can behave like two different oxidants. Shaik used density functional theory to conclude "... that sulfoxidation and N-dealkylation proceed largely via different spin states of Cpd 1," consistent with the two oxidations occurring by different pathways (Sharma et al., 2003). Furthermore, Shaik and coworkers (Ogliaro et al., 2002) used density functional theory to conclude that Cpd 0 is a weak oxidant in comparison to Cpd 1. This has been confirmed with model systems (Park et al., 2006). In contrast, Bach (Bach and Dmitrenko, 2006) used theoretical calculations to conclude that the peroxy species was the active oxidant.

Given the controversy, and the conflicting data, a simple experiment is required to determine whether, and when, multiple oxidants are catalyzing a reaction. One such experiment involves the use of isotopically sensitive branching, which can be used to determine whether products come from rapidly interchanging enzyme-substrate complexes, or whether they arise from kinetically independent enzyme-substrate complexes (Jones et al., 1986; Atkins and Sligar, 1988; Wagschal et al., 1991). Three things led us to consider the possibility that N-oxygenation may occur through Cpd 0. 1) Sulfoxi-

This research was supported by National Institute of Environmental and Health Sciences Grant 09122.

<sup>1</sup> Current affiliation (T.S.D.): Department of Medicinal Chemistry and Pharmacognosy, University of Illinois at Chicago.

Article, publication date, and citation information can be found at <http://dmd.aspetjournals.org>.

doi:10.1124/dmd.106.010280.

**ABBREVIATIONS:** P450, cytochrome P450; Cpd, compound; LC/MS, liquid chromatography/mass spectroscopy.

ation, a reaction closely related to *N*-oxygenation, has been shown by isotopically sensitive branching experiments to occur from a second oxidant (Volz et al., 2002). 2) *N*-Oxygenation possesses a relatively low energy barrier in comparison to aromatic hydroxylation (Dowers et al., 2004), and low barrier reactions have a higher probability of occurring through a reaction with Cpd 0, given its low reactivity. (Ogliaro et al., 2002). 3) The possibility that *N*-oxygenation occurs through Cpd 0 has been suggested in the past (Coon et al., 1998). Therefore, we aimed to test whether *N*-oxygenation occurs through the same active iron-oxygen species as *O*-dealkylation, a P450-mediated reaction that has a relatively high energy barrier in comparison to *N*-dealkylation (Jones et al., 2002) and is generally accepted to occur through Cpd 1 (Lindsay-Smith and Sleath, 1983; Harada et al., 1984; Vaz and Coon, 1994; Higgins et al., 2001; Meunier et al., 2004).


### Materials and Methods

All chemicals were purchased from Sigma-Aldrich (St. Louis, MO), unless otherwise noted. The aromatic hydroxylation product, 6-methoxy-3-quinolinol, was purchased from ChemBridge Corporation (San Diego, CA).  $^2H_3$ -6-Methoxyquinoline was synthesized from 6-hydroxyquinoline and  $^2H_3$ -methyl iodide, which was purchased from Cambridge Isotope Laboratories (Andover, MA). The product was purified on a gravity silica column and verified by liquid chromatography/mass spectroscopy (LC/MS). The deuterium content from the LC/MS was consistent with that in the  $^2H_3$ -methyl iodide (99.5%), and no attempt was made to correct for incomplete incorporation. Incubations were performed using RECO system CYP1A2, purchased from Invitrogen (Carlsbad, CA). Invitrogen's CYP1A2 enzyme mix consists of 0.5  $\mu$ M CYP1A2, 0.2  $\mu$ M NADPH P450 reductase, 0.5  $\mu$ g/ $\mu$ l 3-[(3-cholamidopropyl)dimethylammonio]propanesulfonate, 0.1  $\mu$ g/ $\mu$ l liposomes [dilauroyl phosphatidylcholine, dioleoyl phosphatidylcholine, dilauroyl phosphatidylserine (1:1:1)], 3 mM reduced glutathione, and 50 mM HEPES/KOH, pH 7.4. Invitrogen's buffer mix is in 1 M potassium/sodium phosphate (pH 7.4). The incubations consisted of 200  $\mu$ l enzyme mix, 270  $\mu$ l buffer mix, 1 mM NADPH, and 0.04 mg/ml catalase. Varying concentrations of substrate in methanol were added to the incubations in 5- $\mu$ l quantities. The samples were incubated in a shaker bath for 30 min at 37°C. The reactions were quenched with 3 ml of ethyl acetate, and 9.2 nmol of quinoline *N*-oxide was added as an internal standard. The samples were extracted with ethyl acetate, which was evaporated under nitrogen. The resulting residue was dissolved in approximately 200  $\mu$ l of methanol for LC/MS analysis. LC/MS analysis was performed as described previously (Dowers et al., 2004).

### Results and Discussion

The use of 6-methoxyquinoline as a substrate for P4501A2 allowed us to determine whether aromatic hydroxylation, *N*-oxidation, and *O*-dealkylation come from a common active-oxygen species (Scheme 1). Because isotopically sensitive branching from *O*-demethylation to aromatic hydroxylation has been shown previously (Lindsay-Smith and Sleath, 1983; Harada et al., 1984), we expected branching to occur to the aromatic hydroxylation metabolite of 6-methoxyquinoline (4), but not necessarily to the *N*-oxide (3). Obtaining an inverse isotope effect on the formation of the phenol 4 would ensure that the substrate's motion in the active site is fast relative to the oxidation reaction. In an elegant study by Regal et al. (2005) on the kinetics of caffeine metabolism by CYP1A2, fast interchange was observed for caffeine *N*-dealkylation; thus, fast interchange is likely for 6-methoxyquinoline as well.

Isotopically sensitive branching experiments test whether enzyme-substrate complexes for different products for a single substrate can rapidly interchange (Jones et al., 1986). If interchange is rapid, isotopic substitution will result in branching from the product with an isotope effect to a product that has no isotope effect associated with its formation. If the enzyme-substrate complexes are slow to interchange, no branching will be observed. In the case of 6-methoxyquinoline,



SCHEME 1. Metabolism of 6-methoxyquinoline (1) by P4501A2 to produce 6-hydroxyquinoline (2), 6-methoxyquinoline *N*-oxide (3), and 6-methoxy-3-quinolinol (4).

three products can be formed, as shown in Scheme 1. Substitution of deuterium on the methoxy group should lead to a significant normal isotope effect on the *O*-dealkylation pathway, leading to lower product formation. If aromatic oxidation and *N*-oxide formation occur from enzyme-substrate complexes that are rapidly interchangeable with the *O*-dealkylation enzyme-substrate complex, an inverse isotope effect will be observed on the aromatic oxidation and *N*-oxide products. The overall magnitude of the inverse isotope effect depends on the amount of each product formed and the magnitude of the isotope effect for *O*-dealkylation. If aromatic oxidation or *N*-oxide formation occur from enzyme-substrate complexes that *cannot* interchange with the *O*-dealkylation enzyme-substrate complex, no isotope effect will be observed on the aromatic oxidation or *N*-oxide products. The latter result was observed for *N*-dealkylation and sulfoxidation (Volz et al., 2002) of *N,N*-dimethyl-4-(methylthio)aniline, indicating that these two products do not come from interchangeable enzyme-substrate complexes and most likely come from two different oxidants. Given previous experiments that indicated that aromatic oxidation occurred from rapidly interchanging enzyme-substrate complexes, we anticipated two possible outcomes upon deuteration of the methoxy group: 1) a decrease in 2 as a result of a normal isotope effect, an increase in aromatic oxidation to form 4, and the same amount of *N*-oxide formation, or 2) a decrease in product 2 and an increase in products 3 and 4 as a result of branching. Situation 1 would implicate a second oxidant for *N*-oxide formation, and situation 2 would indicate a common oxidant for all three products.

Large inverse isotope effects were observed on both *N*-oxygenation and aromatic hydroxylation of 6-methoxyquinoline (Table 1). The results are only consistent with all three products arising from a kinetically indistinguishable (rapidly interchanging) enzyme-substrate complex, i.e., the same oxidant. However, some fraction of the products may still come from a second noninterchangeable enzyme-substrate complex. If this is the case, only a fraction of the intrinsic isotope effect would be observed.

Because *N*-oxide formation and aromatic oxidation are minor pathways, the majority of the isotope effect is observed on these pathways, with only a small isotope effect on *O*-dealkylation. The lower limit of the intrinsic isotope effect can be estimated to be 6.5 by dividing the isotope effect on *O*-dealkylation (1.3) by the inverse isotope effect for either *N*-oxide formation (0.20) or the aromatic oxidation product (0.17). Some of the observed intramolecular and branched isotope effects for P450 *O*-dealkylation reactions that have been reported are

TABLE 1  
Isotope effects for metabolism of protio and deuterio 6-methoxyquinoline (1)

	6-OH-Quinoline 2	6-Methoxyquinoline N-Oxide 3	6-Methoxy-3-Quinololinol 4
$(V/K)_H^a$	1994 ± 138	9.1 ± 1.1	24 ± 1.1
$(V/K)_D$	1552 ± 151	45 ± 15	140 ± 26
$(V/K)_H/(V/K)_D$	1.3 ± 0.22	0.20 ± 0.09	0.17 ± 0.04

<sup>a</sup> The  $V_{max}$  and  $K_m$  values were obtained by plotting the rate of product formation versus substrate concentration. The data were fit to Michaelis-Menton kinetics using the program EnzFitter. At higher substrate concentrations, a change in the ratio of product 3 to product 4 was observed, indicating two substrates binding in the active site. Therefore, the  $V_{max}$  and  $K_m$  values were obtained at low substrate concentrations (3  $\mu$ M to 50  $\mu$ M). Possible errors associated with two substrates binding should affect the  $V_{max}$  and  $K_m$  values of the protio and deuterio substrates equally and, therefore, would have no effect on  $(V/K)_H/(V/K)_D$ .

7.3 for anisole (Lindsay-Smith and Sleath, 1983), 5.1 (Watanabe et al., 1982) and 8.4 (Guengerich et al., 2004) for 4-methoxyanisole, 7.0 for methacetin (Guengerich et al., 2004), 8.4 for 4-cyanoanisole (Guengerich et al., 2004), 14.7 for 4-nitroanisole, and 5.5 for 7-ethoxycoumarin (Miwa et al., 1985). Thus, the magnitude of this isotope effect is consistent with other isotope effects that have been measured for P450-mediated O-dealkylation reactions indicating that the majority, or all, of the product comes from rapidly interchangeable enzyme-substrate complexes and, therefore, from the same oxidant. Again, however, this does not prove that 100% of the N-oxide arises from a single oxidant.

The strongest evidence that all the products come from the same oxidant is that the amount of branching to the N-oxide (3) and 6-methoxy-3-quinolinol (4) is identical. Isotopically sensitive branching requires that the relative amount of branching from a major pathway to alternate products will be the same for any two products; i.e., that the inverse isotope effects be the same for both products if the enzyme-substrate complexes are in rapid equilibrium (Korzekwa et al., 1995). However, if two oxidants catalyze the reactions that occur to form 3 and 4, the fraction of 3 and 4 that comes from a given oxidant will most likely be different, reflecting the different activation energies for the two reactions. If this is the case, the observed branching isotope effects would be different, with a smaller inverse isotope effect for the pathway that favors the second oxidant. Clearly, this not the result obtained (see Table 1). Thus, for both the reactions to occur from both oxidants, it is required that the relative energy of formation of 3 and 4 be identical for both oxidizing species. In fact, mutants that purportedly enhance Cpd 0 have been shown to alter product ratios and rates of reaction in every case, indicating a difference in the energetics of the two oxidants (Vaz et al., 1998; Vatsis and Coon, 2002; Volz et al., 2002; Jin et al., 2003; Chandrasena et al., 2004) and consistent with computational results (Ogliaro et al., 2002).

From these results it is apparent that for the three reactions in Scheme 1 a single oxidant is responsible for most, if not all, of the products formed. The nature of the oxidant remains a matter of debate. However, the consensus oxidant for O-dealkylation reactions is Cpd 1. Since the results herein require a single oxidant for three energetically different reactions, it is most likely the iron-oxo species Cpd 1.

Department of Chemistry  
Washington State University  
Pullman, Washington

TAMARA S. DOWERS<sup>1</sup>  
JEFFREY P. JONES

## References

- Atkins WM and Sligar SG (1988) Deuterium isotope effects in norcamphor metabolism by cytochrome P-450cam: kinetic evidence for the two-electron reduction of a high-valent iron-oxo intermediate. *Biochemistry* **27**:1610–1616.
- Bach RD and Dmitrenko O (2006) The "somersault" mechanism for the P-450 hydroxylation of hydrocarbons. The invention of transient inverted metastable hydroperoxides. *J Am Chem Soc* **128**:1474–1488.
- Chandrasena REP, Vatsis KP, Coon MJ, Hollenberg PF, and Newcomb M (2004) Hydroxylation by the hydroperoxy-iron species in cytochrome P450 enzymes. *J Am Chem Soc* **126**:115–126.
- Coon MJ, Vaz AD, McGinnity DF, and Peng HM (1998) Multiple activated oxygen species in P450 catalysis: contributions to specificity in drug metabolism. *Drug Metab Dispos* **26**:1190–1193.

- Dowers TS, Rock DA, Perkins BNS, and Jones JP (2004) An analysis of the regioselectivity of aromatic hydroxylation and N-oxygenation by cytochrome P450 enzymes. *Drug Metab Dispos* **32**:328–332.
- Groves JT, McClusky GA, White RE, and Coon MJ (1978) Aliphatic hydroxylation by highly purified liver microsomal cytochrome P450. Evidence for a carbon radical intermediate. *Biochem Biophys Res Commun* **81**:154–160.
- Guengerich FP, Krauser JA, and Johnson WW (2004) Rate-limiting steps in oxidations catalyzed by rabbit cytochrome P450 1A2. *Biochemistry* **43**:10775–10788.
- Guengerich FP and MacDonald TL (1990) Mechanisms of cytochrome P-450 catalysis. *FASEB J* **4**:2453–2459.
- Harada N, Miwa GT, Walsh JS, and Lu AYH (1984) Kinetic isotope effects on cytochrome P450-catalyzed oxidation reactions: evidence for the irreversible formation of an activated oxygen intermediate of cytochrome P-448. *J Biol Chem* **259**:3005–3010.
- Higgins L, Korzekwa KR, Rao S, Shou M, and Jones JP (2001) An assessment of the reaction energetics for cytochrome P450-mediated reactions. *Arch Biochem Biophys* **385**:220–230.
- Hutzler JM, Powers FJ, Wynalda MA, and Wienkers LC (2003) Effect of carbonate anion on cytochrome P450 2D6-mediated metabolism in vitro: the potential role of multiple oxygenating species. *Arch Biochem Biophys* **417**:165–175.
- Jin S, Makris TM, Bryson TA, Sligar SG, and Dawson JH (2003) Epoxidation of olefins by hydroperoxy-ferric cytochrome P450. *J Am Chem Soc* **125**:3406–3407.
- Jones JP, Korzekwa KR, Rettie AE, and Trager WF (1986) Isotopically sensitive branching and its effect on the observed intramolecular isotope effects in cytochrome P450 catalyzed reactions: a new method for the estimation of intrinsic isotope effects. *J Am Chem Soc* **108**:7074–7078.
- Jones JP, Mysinger M, and Korzekwa KR (2002) Computational models for cytochrome P450: a predictive electronic model for aromatic oxidation and hydrogen atom abstraction. *Drug Metab Dispos* **30**:7–12.
- Korzekwa KR, Gillette JR, and Trager WF (1995) Isotope effect studies on the cytochrome P450 enzymes. *Drug Metab Rev* **27**:45–59.
- Lindsay-Smith JR and Sleath PR (1983) Model systems for cytochrome P450 dependent mono-oxygenases. Part 2. Kinetic isotope effects for the oxidative demethylation of anisole and [Me-<sup>3</sup>H]<sub>2</sub>anisole by cytochrome P450 dependent mono-oxygenases and model systems. *J Chem Soc Perkin Trans II* 621–628.
- Meunier B, de Visser SP, and Shaik S (2004) Mechanism of oxidation reactions catalyzed by cytochrome P450 enzymes. *Chem Rev* **104**:3947–3980.
- Miwa GT, Harada N, and Lu AY (1985) Kinetic isotope effects on cytochrome P-450-catalyzed oxidation reactions: full expression of the intrinsic isotope effect during the O-deethylation of 7-ethoxycoumarin by liver microsomes from 3-methylcholanthrene-induced hamsters. *Arch Biochem Biophys* **239**:155–162.
- Ogliaro F, de Visser SP, Cohen S, Sharma PK, and Shaik S (2002) Searching for the second oxidant in the catalytic cycle of cytochrome P450: a theoretical investigation of the iron(III)-hydroperoxy species and its epoxidation pathways. *J Am Chem Soc* **124**:2806–2817.
- Ortiz de Montellano PR (1995) *Cytochrome P450: Structure, Mechanism, and Biochemistry*. Plenum, New York.
- Park MJ, Lee J, Suh Y, Kim J, and Nam W (2006) Reactivities of mononuclear non-heme iron intermediates including evidence that iron(III)-hydroperoxy species is a sluggish oxidant. *J Am Chem Soc* **128**:2630–2634.
- Regal KA, Kunze KL, Peter RM, and Nelson SD (2005) Oxidation of caffeine by CYP1A2: isotope effects and metabolic switching. *Drug Metab Dispos* **33**:1837–1844.
- Sharma PK, De Visser SP, and Shaik S (2003) Can a single oxidant with two spin states masquerade as two different oxidants? A study of the sulfoxidation mechanism by cytochrome p450. *J Am Chem Soc* **125**:8698–8699.
- Vatsis KP and Coon MJ (2002) Ipso-substitution by cytochrome P450 with conversion of p-hydroxybenzene derivatives to hydroquinone: evidence for hydroperoxy-iron as the active oxygen species. *Arch Biochem Biophys* **397**:119–129.
- Vaz AD and Coon MJ (1994) On the mechanism of action of cytochrome P450: evaluation of hydrogen abstraction in oxygen-dependent alcohol oxidation. *Biochemistry* **33**:6442–6449.
- Vaz AD, McGinnity DF, and Coon MJ (1998) Epoxidation of olefins by cytochrome P450: evidence from site-specific mutagenesis for hydroperoxy-iron as an electrophilic oxidant. *Proc Natl Acad Sci USA* **95**:3555–3560.
- Volz TJ, Rock DA, and Jones JP (2002) Evidence for two different active oxygen species in cytochrome P450 BM3 mediated sulfoxidation and N-dealkylation reactions. *J Am Chem Soc* **124**:9724–9725.
- Wagschal K, Savage TJ, and Croteau R (1991) Isotopically sensitive branching as a tool for evaluating multiple product formation by monoterpene cyclases. *Tetrahedron* **47**:5933–5944.
- Watanabe Y, Shigeru O, and Iyanagi T (1982) Mechanism of enzymatic S-oxygenation of thioanisole derivative and O-demethylation of anisole derivatives. *Bull Chem Soc Jpn* **55**: 188–195.

Address correspondence to: Jeffrey P. Jones, Department of Chemistry, 477 Fulmer Synthesis, Washington State University, Pullman, WA 99164. E-mail jpp@wsu.edu