

> [Dermatol Surg.](#) 2014 Oct;40(10):1094-102. doi: 10.1097/01.DSS.0000452662.69539.57.

Temperature-modulated photodynamic therapy for the treatment of actinic keratosis on the extremities: a pilot study

Andrea Willey¹, R Rox Anderson, Fernanda H Sakamoto

Affiliations + expand

PMID: 25207759 DOI: 10.1097/01.DSS.0000452662.69539.57

Abstract

Background: The efficacy of photodynamic therapy (PDT) using topical 5-aminolevulinic acid (ALA) for the treatment of actinic keratosis (AKs) is lower on the distal extremities compared with the head and neck areas. The strong temperature dependence of porphyrin synthesis in biologic tissue suggests that heating skin during incubation may improve the efficacy of PDT, particularly in areas where biologic temperatures are naturally lower. The aim of this study was to evaluate the efficacy and tolerability of temperature-modulated PDT for the treatment of AKs on the extremities.

Methods: In this IRB-approved, single-center study, the upper or lower extremities of 20 subjects were treated with 20% ALA under occlusion, followed by 10 J/cm, 417-nm blue light. One of the 2 extremities treated was heated during the 1-hour incubation. Outcome measures included lesion counts, tolerability, and global improvement at baseline, 1 week, and 2 and 6 months after treatment.

Results: The median temperatures of the heated and control sides were 38.8°C and 29.4°C, respectively. The median clearance for the heated side was significantly greater than the control side at 2 and 6 months ($p < .0001$). Typical PDT side effects were greater on the heated side compared with the control yet were well tolerated by all subjects.

Conclusion: Warming the skin during incubation of ALA seems to improve the efficacy of PDT in the treatment of AKs on the extremities and is well tolerated when heat application is controlled within the limits of safety.

Similar articles

[Temperature-Modulated Photodynamic Therapy for the Treatment of Actinic Keratosis on the Extremities: A One-Year Follow-up Study.](#)

Willey A, Anderson RR, Sakamoto FH.

Dermatol Surg. 2015 Nov;41(11):1290-5. doi: 10.1097/DSS.0000000000000512.

PMID: 26445287

[A multicenter, randomized, vehicle-controlled phase 2 study of blue light photodynamic therapy with aminolevulinic acid HCl 20% topical solution for the treatment of actinic keratoses on the upper extremities: the effect of occlusion during the drug incubation period.](#)

Schmieder GJ, Huang EY, Jarratt M.

J Drugs Dermatol. 2012 Dec;11(12):1483-9.

PMID: 23377520 Clinical Trial.

[A randomized, blinded, bilateral intraindividual, vehicle-controlled photodynamic therapy with 5-aminolevulinic acid and blue light](#)

FULL TEXT LINKS



ACTIONS



SHARE



PAGE NAVIGATION

Title & authors

< Abstract

Similar articles

Cited by

MeSH terms

Substances

Related information

LinkOut - more resources

Biofrontera Exhibit 1032
Biofrontera Inc. et al. v. DUSA Pharmaceuticals, Inc.
IPR2022-00056

> *Dermatol Surg.* 2014 Oct;40(10):1094-102. doi: 10.1097/01.DSS.0000452662.69539.57.

Temperature-modulated photodynamic therapy for the treatment of actinic keratosis on the extremities: a pilot study

Andrea Willey¹, R Rox Anderson, Fernanda H Sakamoto

Affiliations + expand

PMID: 25207759 DOI: 10.1097/01.DSS.0000452662.69539.57

Abstract

Background: The efficacy of photodynamic therapy (PDT) using topical 5-aminolevulinic acid (ALA) for the treatment of actinic keratosis (AKs) is lower on the distal extremities compared with the head and neck areas. The strong temperature dependence of porphyrin synthesis in biologic tissue suggests that heating skin during incubation may improve the efficacy of PDT, particularly in areas where biologic temperatures are naturally lower. The aim of this study was to evaluate the efficacy and tolerability of temperature-modulated PDT for the treatment of AKs on the extremities.

Methods: In this IRB-approved, single-center study, the upper or lower extremities of 20 subjects were treated with 20% ALA under occlusion, followed by 10 J/cm, 417-nm blue light. One of the 2 extremities treated was heated during the 1-hour incubation. Outcome measures included lesion counts, tolerability, and global improvement at baseline, 1 week, and 2 and 6 months after treatment.

Results: The median temperatures of the heated and control sides were 38.8°C and 29.4°C, respectively. The median clearance for the heated side was significantly greater than the control side at 2 and 6 months ($p < .0001$). Typical PDT side effects were greater on the heated side compared with the control yet were well tolerated by all subjects.

Conclusion: Warming the skin during incubation of ALA seems to improve the efficacy of PDT in the treatment of AKs on the extremities and is well tolerated when heat application is controlled within the limits of safety.

Similar articles

Temperature-Modulated Photodynamic Therapy for the Treatment of Actinic Keratosis on the Extremities: A One-Year Follow-up Study.

Willey A, Anderson RR, Sakamoto FH.

Dermatol Surg. 2015 Nov;41(11):1290-5. doi: 10.1097/DSS.0000000000000512.

PMID: 26445287

A multicenter, randomized, vehicle-controlled phase 2 study of blue light photodynamic therapy with aminolevulinic acid HCl 20% topical solution for the treatment of actinic keratoses on the upper extremities: the effect of occlusion during the drug incubation period.

Schmieder GJ, Huang EY, Jarratt M.

J Drugs Dermatol. 2012 Dec;11(12):1483-9.

PMID: 23377520 Clinical Trial.

A randomized, blinded, bilateral intraindividual, vehicle-controlled trial of the use of photodynamic therapy with 5-aminolevulinic acid and blue light for the treatment of actinic keratosis on the face.

FULL TEXT LINKS



ACTIONS



SHARE



PAGE NAVIGATION

Title & authors

< Abstract

Similar articles

Cited by

MeSH terms

Substances

Related information

LinkOut - more resources

J Drugs Dermatol. 2011 Sep;10(9):1049-56.
PMID: 22052276 Clinical Trial.

Spanish-Portuguese consensus statement on use of daylight-mediated photodynamic therapy with methyl aminolevulinic acid in the treatment of actinic keratosis.

Gilaberte Y, Aguilar M, Almagro M, Correia O, Guillén C, Harto A, Pérez-García B, Pérez-Pérez L, Redondo P, Sánchez-Carpintero I, Serra-Guillén C, Valladares LM.
Actas Dermosifiliogr. 2015 Oct;106(8):623-31. doi: 10.1016/j.ad.2015.06.001. Epub 2015 Jun 24.
PMID: 26115793 English, Spanish.

Current treatments of actinic keratosis.

Gold MH, Nestor MS.
J Drugs Dermatol. 2006 Feb;5(2 Suppl):17-25.
PMID: 16485877 Review.

[See all similar articles](#)

Cited by 8 articles

Comparison of Blue and White Lamp Light with Sunlight for Daylight-Mediated, 5-ALA Photodynamic Therapy, in vivo.

Marra K, LaRochelle EP, Chapman MS, Hoopes PJ, Lukovits K, Maytin EV, Hasan T, Pogue BW.
Photochem Photobiol. 2018 Sep;94(5):1049-1057. doi: 10.1111/php.12923. Epub 2018 May 16.
PMID: 29663426 [Free PMC article](#).

ALA-PpIX mediated photodynamic therapy of malignant gliomas augmented by hypothermia.

Fisher CJ, Niu C, Foltz W, Chen Y, Sidorova-Darmos E, Eubanks JH, Lilge L.
PLoS One. 2017 Jul 31;12(7):e0181654. doi: 10.1371/journal.pone.0181654. eCollection 2017.
PMID: 28759636 [Free PMC article](#).

Photodynamic therapy in dermatology beyond non-melanoma cancer: An update.

Wen X, Li Y, Hamblin MR.
Photodiagnosis Photodyn Ther. 2017 Sep;19:140-152. doi: 10.1016/j.pdpdt.2017.06.010. Epub 2017 Jun 21.
PMID: 28647616 [Free PMC article](#). Review.

Effect of Expedited Microneedle-Assisted Photodynamic Therapy for Field Treatment of Actinic Keratoses: A Randomized Clinical Trial.

Petukhova TA, Hassoun LA, Foolad N, Barath M, Sivamani RK.
JAMA Dermatol. 2017 Jul 1;153(7):637-643. doi: 10.1001/jamadermatol.2017.0849.
PMID: 28514458 [Free PMC article](#). Clinical Trial.

Current Advances in 5-Aminolevulinic Acid Mediated Photodynamic Therapy.

Thunshelle C, Yin R, Chen Q, Hamblin MR.
Curr Dermatol Rep. 2016 Sep;5(3):179-190. doi: 10.1007/s13671-016-0154-5. Epub 2016 Jul 13.
PMID: 28163981 [Free PMC article](#).

[See all "Cited by" articles](#)

MeSH terms

- > Aged
- > Aged, 80 and over
- > Aminolevulinic Acid / therapeutic use*
- > Blister / etiology
- > Clinical Protocols
- > Erythema / etiology
- > Extremities
- > Female
- > Humans
- > Keratosis, Actinic / drug therapy*
- > Male
- > Middle Aged
- > Pain / etiology
- > Photochemotherapy / adverse effects
- > Photochemotherapy / methods*
- > Photosensitizing Agents / therapeutic use*
- > Pilot Projects

[> Treatment Outcome](#)

Substances

- [> Photosensitizing Agents](#)
- [> Aminolevulinic Acid](#)

Related information

[MedGen](#)

[PubChem Compound \(MeSH Keyword\)](#)

LinkOut - more resources

Full Text Sources

[Ovid Technologies, Inc.](#)

[Wiley](#)

[Wolters Kluwer](#)

Other Literature Sources

[The Lens - Patent Citations](#)

[scite Smart Citations](#)

[NCBI Literature Resources](#) [MeSH](#) [PMC](#) [Bookshelf](#) [Disclaimer](#)

FOLLOW NCBI



Follow NLM



National Library of Medicine
8600 Rockville Pike
Bethesda, MD 20894

Copyright
FOIA
Privacy

Help
Accessibility
Careers

NLM | NIH | HHS | USA.gov

Cited In for PMID: 25207759

9 results

- 1 **Temperature-modulated photodynamic therapy for the treatment of actinic keratosis on the extremities: a pilot study.**
Willey A, Anderson RR, Sakamoto FH.
Dermatol Surg. 2014 Oct;40(10):1094-102. doi: 10.1097/01.DSS.0000452662.69539.57.
PMID: 25207759
- 2 **Comparison of Blue and White Lamp Light with Sunlight for Daylight-Mediated, 5-ALA Photodynamic Therapy, in vivo.**
Marra K, LaRochelle EP, Chapman MS, Hoopes PJ, Lukovits K, Maytin EV, Hasan T, Pogue BW.
Photochem Photobiol. 2018 Sep;94(5):1049-1057. doi: 10.1111/php.12923. Epub 2018 May 16.
PMID: 29663426 [Free PMC article.](#)
- 3 **ALA-PpIX mediated photodynamic therapy of malignant gliomas augmented by hypothermia.**
Fisher CJ, Niu C, Foltz W, Chen Y, Sidorova-Darmos E, Eubanks JH, Lilge L.
PLoS One. 2017 Jul 31;12(7):e0181654. doi: 10.1371/journal.pone.0181654. eCollection 2017.
PMID: 28759636 [Free PMC article.](#)
- 4 **Photodynamic therapy in dermatology beyond non-melanoma cancer: An update.**
Wen X, Li Y, Hamblin MR.
Photodiagnosis Photodyn Ther. 2017 Sep;19:140-152. doi: 10.1016/j.pdpdt.2017.06.010. Epub 2017 Jun 21.
PMID: 28647616 [Free PMC article.](#) Review.
- 5 **Effect of Expedited Microneedle-Assisted Photodynamic Therapy for Field Treatment of Actinic Keratoses: A Randomized Clinical Trial.**
Petukhova TA, Hassoun LA, Foolad N, Barath M, Sivamani RK.
JAMA Dermatol. 2017 Jul 1;153(7):637-643. doi: 10.1001/jamadermatol.2017.0849.
PMID: 28514458 [Free PMC article.](#) Clinical Trial.
- 6 **Current Advances in 5-Aminolevulinic Acid Mediated Photodynamic Therapy.**
Thunshelle C, Yin R, Chen Q, Hamblin MR.
Curr Dermatol Rep. 2016 Sep;5(3):179-190. doi: 10.1007/s13671-016-0154-5. Epub 2016 Jul 13.
PMID: 28163981 [Free PMC article.](#)
- 7 **In-office Painless Aminolevulinic Acid Photodynamic Therapy: A Proof of Concept Study and Clinical Experience in More Than 100 Patients.**
Martin GM.
J Clin Aesthet Dermatol. 2016 Feb;9(2):19-26.
PMID: 27047629 [Free PMC article.](#)
- 8 **Comprehensive analytical model for CW laser induced heat in turbid media.**
Erkol H, Nouizi F, Luk A, Unlu MB, Gulsen G.
Opt Express. 2015 Nov 30;23(24):31069-84. doi: 10.1364/OE.23.031069.
PMID: 26698736 [Free PMC article.](#)
- 9 **Pre-treatment protoporphyrin IX concentration in actinic keratosis lesions may be a predictive biomarker of response to aminolevulinic-acid based photodynamic therapy.**
Kanick SC, Davis SC, Zhao Y, Sheehan KL, Hasan T, Maytin EV, Pogue BW, Chapman MS.
Photodiagnosis Photodyn Ther. 2015 Dec;12(4):561-6. doi: 10.1016/j.pdpdt.2015.10.006. Epub 2015 Oct 22.
PMID: 26480810 [Free PMC article.](#) Clinical Trial.

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.